arXiv:1212.0990v1 [hep-ph] 5 Dec 2012

The Effective Chiral Lagrangian From Dimension-Six Parity and Time-Reversal Violation

J. de Vries[1,2], **E. Mereghetti**[3], **R. G. E. Timmermans**[1], **and U. van Kolck**[4,5]

[1] *KVI, Theory Group, University of Groningen,*
9747 AA Groningen, The Netherlands

[2] *Nikhef, Science Park 105,*
1098 XG Amsterdam, The Netherlands

[3] *Ernest Orlando Lawrence Berkeley National Laboratory, University of California,*
Berkeley, CA 94720, USA

[4] *Institut de Physique Nucléaire, Université Paris Sud, CNRS/IN2P3,*
91406 Orsay, France

[5] *Department of Physics, University of Arizona,*
Tucson, AZ 85721, USA

Abstract

We classify the parity- and time-reversal-violating operators involving quark and gluon fields that have effective dimension six: the quark electric dipole moment, the quark and gluon chromo-electric dipole moments, and four four-quark operators. We construct the effective chiral Lagrangian with hadronic and electromagnetic interactions that originate from them and serves as the basis for calculations of low-energy observables. The form of the effective interactions depends on the chiral properties of these operators. We develop a power-counting scheme and calculate within this scheme, as an example, the parity- and time-reversal-violating pion-nucleon form factor. We also discuss the electric dipole moments of the nucleon and light nuclei.

1 Introduction

The Standard Model (SM) of particle physics contains, in its minimal version, two sources of time-reversal (T), or, equivalently, CP violation. In the electroweak sector, the phase in the quark mixing matrix [1] is associated with the Jarlskog parameter $J_{\rm CP} \simeq 3 \times 10^{-5}$ [2]. The strong sector contains the QCD vacuum angle $\bar{\theta}$ [3], but the experimental upper limit on the neutron electric dipole moment (EDM) shows that $\bar{\theta}$ is unnaturally small, $\bar{\theta} \lesssim 10^{-10}$ [4]. CP violation within the SM is believed to be insufficient for a successful baryogenesis scenario [5], and therefore new sources of CP violation are expected in order to explain the cosmological matter-antimatter asymmetry. This is not a surprise since the SM is likely but the dimension-four part of an effective field theory (EFT) that contains higher-dimensional operators, some of which will violate CP.

Powerful probes of such CP violation beyond the SM are EDMs of nucleons, nuclei, atoms, and molecules [6, 7], which violate both parity and time reversal ($\not{P}\not{T}$). Since the SM predictions from the quark mixing matrix [8] are orders of magnitude away from current experimental limits, a finite EDM in upcoming experiments would be an unambiguous sign of new physics. In addition to impressive improvements [9] on the time-honored EDM experiments with neutrons and neutral atoms, in particular ^{199}Hg, which have resulted in very precise limits [10, 11], novel ideas exist for the measurement of EDMs of charged particles, such as the proton, deuteron and helion, in storage rings [12]. An important question that comes up is whether, when future experiments measure nonzero EDMs, we will be able to pinpoint the microscopic source of P and T violation.

EDMs of strongly interacting particles arise from the higher-dimensional $\not{P}\not{T}$ operators at the quark-gluon level. These non-renormalizable operators might have their origin in a renormalizable theory at a higher-energy scale, such as, for example, supersymmetric (SUSY) extensions of the SM. At the SM scale, the most important higher-dimensional $\not{P}\not{T}$ operators should be those of dimension six, as we are not concerned here with CP violation in the dimension-five leptonic operator [13] that gives rise to neutrino masses and mixings. From symmetry considerations it is found [14, 15, 16, 17] that the following flavor-diagonal $\not{P}\not{T}$ operators appear at an effective dimension six: the quark electric dipole moment (qEDM) [18], which couples quarks and photons; the quark chromo-electric dipole moment (qCEDM) [19], which couples quarks and gluons; the Weinberg operator [20], which couples three gluons and gives rise to a gluon chromo-electric dipole moment (gCEDM) [21], and four four-quark operators [16, 22].

Since it is not feasible to calculate hadronic and nuclear properties directly from a Lagrangian at the quark and gluon level, we use chiral EFT [23] —a generalization to more than one nucleon of chiral perturbation theory (χPT) [24]— to translate microscopic operators into operators that include nucleons, pions, and photons. (For reviews, see Refs. [25, 26, 27].) After the translation, we are able to calculate hadronic properties directly from the effective Lagrangian. For the dimension-four $\bar{\theta}$ term, this method was first employed in Ref. [4], and later extended in the context of $SU(2)\times SU(2)$ [28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38] and of $SU(3)\times SU(3)$ [39, 40] χPT.

In this paper we extend the method further to include dimension-six operators in the framework of $SU(2) \times SU(2)$ χPT. (Generalization to $SU(3) \times SU(3)$ is straightforward.) Different $\not{P}\not{T}$ sources at the quark-gluon level produce different hadronic interactions. The effective chiral Lagrangian includes not only interactions that stem from spontaneous chiral symmetry breaking and are therefore chiral invariant, but also interactions that break chiral symmetry in the same way as chiral-symmetry-breaking operators at the QCD level. Since the dimension-six operators break chiral symmetry differently from each other and from the $\bar{\theta}$ term, they will generate different low-energy hadronic interactions. Given enough observables it should be thus possible to separate the various $\not{P}\not{T}$ sources.

In addition to constructing the Lagrangian, we need to organize in leading order (LO), next-to-leading order (NLO), *etc.* the various effective $\not{P}\not{T}$ operators that appear. This is done according to the estimated size of their contributions to observables. In order to get a consistent, manifest power counting we work in the heavy-baryon framework [41] wherein the nucleon mass has been eliminated from the nucleon propagator. This framework has a transparent power counting and greatly simplifies loop calculations, but there are some complications when one goes to subleading orders in the Lagrangian. These problems can be solved by demanding that the Lagrangian obeys reparametrization invariance (RPI) [42]. This puts constraints on certain coefficients of operators, which we construct up to NNLO.

In Refs. [43, 33, 34, 35, 36] we have used some of the effective interactions to calculate the EDMs of the nucleons and the lightest nuclei. It was found that in LO they depend on six low-energy constants (LECs), which for each fundamental $\not{P}\not{T}$ source have different relative sizes. The idea is that, from EDM measurements, the LECs can be inferred and the dominant fundamental source identified. For example, if the deuteron EDM is significantly larger than the sum of the nucleon EDMs, this points towards new physics in form of a qCEDM [34, 36] or, as we will demonstrate here, a particular isospin-breaking $\not{P}\not{T}$ four-quark operator. If the deuteron EDM is well approximated by the nucleon EDMs, but the helion (triton) EDM is far away from the neutron (proton) EDM, it is a hint for the SM $\bar{\theta}$ term. The deuteron magnetic quadrupole moment (MQM), if experimentally accessible, could play an important role as well [34, 37].

In this article, we construct the full Lagrangian necessary to perform these and other low-energy $\not{P}\not{T}$ calculations. We follow the approach outlined in Refs. [30, 32], where the χPT Lagrangian originating from the $\bar{\theta}$ term was derived, and the higher-dimensional interactions were examined. We identify, for each source, the size of the six $\not{P}\not{T}$ interactions relevant for light nuclear EDMs. Apart from that, we construct also all other operators that appear at the same order. These operators could play a role in the calculation of other observables that violate P and T such as the EDMs of heavier nuclei or of ions and atoms. $\not{P}\not{T}$ form factors and scattering observables could depend on these operators as well. An important ingredient in calculating hadronic and nuclear $\not{P}\not{T}$ observables is the $\not{P}\not{T}$ pion-nucleon form factor. We investigate this form factor here. Finally, we extend our earlier light-nuclear EDM calculations to a four-quark operator that has recently been shown to arise below the SM scale from weak-boson exchange [22].

Our paper is organized as follows. In Secs. 2 and 3 we discuss the QCD Lagrangian, the $\bar{\theta}$ term, and the possible higher-dimensional $\not{P}\not{T}$ operators. In Sec. 4 we briefly discuss $SU(2) \times SU(2)$ χPT, and how to incorporate the $\not{P}\not{T}$ operators in this framework. The bulk of the article is Secs. 5, 6, and 8, where we construct the χPT Lagrangian up to NNLO including the removal of pion tadpoles that proliferate when $\not{P}\not{T}$ appears together with isospin violation. In Sec. 7 we use the constructed Lagrangian to calculate the $\not{P}\not{T}$ pion-nucleon form factor (PNFF), and in Sec. 9 we give the electric dipole form factor (EDFF) of the nucleon in case of the specific $\not{P}\not{T}$ four-quark operator of Ref. [22]. In Sec. 10 the extra elements that arise in nuclear systems are addressed. We discuss our results and conclude in Sec. 11.

2 The underlying quark-gluon Lagrangian

Before we discuss higher-dimensional $\not{P}\not{T}$ operators, we set our notation by recalling the main SM ingredients. The SM Lagrangian is completely determined by gauge symmetry with gauge group $SU_c(3) \times SU_L(2) \times U_Y(1)$, by the matter content, and by the requirement of renormalizability.

In its minimal form, which we assume in this paper, the matter content consists of three generations of leptons and quarks, and one scalar doublet. Quark and lepton fields carry a generation index $r = 1, 2, 3$, which we will often leave implicit, running over the three generations of up-type quarks $u = (u, c, t)$, down-type quarks $d = (d, s, b)$, charged leptons $e = (e, \mu, \tau)$ and neutrinos $\nu = (\nu_e, \nu_\mu, \nu_\tau)$. The left-handed fermions are doublets of $SU_L(2)$,

$$q_L = \left(\begin{array}{c} u_L \\ d_L \end{array} \right), \qquad l_L = \left(\begin{array}{c} \nu_L \\ e_L \end{array} \right), \tag{1}$$

while the right-handed fields u_R, d_R and e_R are singlets. The field φ denotes an $SU_L(2)$ doublet of scalar fields φ^J, $J = 1, 2$. For convenience we define $\tilde{\varphi}^I = \epsilon^{IJ} \varphi^{J*}$, where ϵ^{IJ} is the antisymmetric tensor in two dimensions ($\epsilon^{12} = +1$). Left and right-handed quarks are in the fundamental representation of $SU_c(3)$, while leptons and scalars are singlets. We sometimes group the right-handed up- and down-type quarks in a doublet q_R, and define quark doublets $q = q_L + q_R$. The hypercharge assignments under the group $U_Y(1)$ are $1/6$, $2/3$, $-1/3$, $-1/2$, -1, and $1/2$ for q_L, u_R, d_R, l_L, e_R, and φ, respectively.

We denote the gauge bosons associated with the gauge groups $SU_c(3)$, $SU_L(2)$, and $U_Y(1)$ by, respectively, G^a_μ, W^i_μ, and B_μ, with $a = 1, \ldots, 8$ and $i = 1, 2, 3$. Gauge invariance is most easily imposed by employing covariant combinations of the gauge and matter fields. The covariant derivative of matter fields is

$$D_\mu = \partial_\mu - i\frac{g_s}{2} G^a_\mu \lambda^a - i\frac{g}{2} W^i_\mu \tau^i - ig' Y B_\mu, \tag{2}$$

where g_s, g, and g' are the $SU_c(3)$, $SU_L(2)$, and $U_Y(1)$ coupling constants; and $\lambda^a/2$ and $\tau^i/2$ are $SU(3)$ and $SU(2)$ generators, in the representation of the field on which the derivative acts. For example, for left-handed quarks λ^a and τ^i are, respectively, the Gell-Mann color and Pauli isospin matrices. The field strengths are

$$\begin{aligned} G^a_{\mu\nu} &= \partial_\mu G^a_\nu - \partial_\mu G^a_\mu - g_s f^{abc} G^b_\mu G^c_\nu, && (3) \\ W^i_{\mu\nu} &= \partial_\mu W^i_\nu - \partial_\nu W^i_\mu - g\epsilon^{ijk} W^j_\mu W^k_\nu, && (4) \\ B_{\mu\nu} &= \partial_\mu B_\nu - \partial_\nu B_\mu, && (5) \end{aligned}$$

with f^{abc} and ϵ^{ijk} denoting the $SU(3)$ and $SU(2)$ structure constants.

The SM Lagrangian is expressed in terms of all possible dimension-four gauge-invariant operators:

$$\begin{aligned} \mathcal{L}_{\rm SM} = & -\frac{1}{4}\left(G^a_{\mu\nu} G^{a\,\mu\nu} + W^i_{\mu\nu} W^{i\,\mu\nu} + B_{\mu\nu} B^{\mu\nu} \right) \\ & + \bar{q}_L i\not{D}\, q_L + \bar{u}_R i\not{D}\, u_R + \bar{d}_R i\not{D}\, d_R + \bar{l}_L i\not{D}\, l_L + \bar{e}_R i\not{D}\, e_R + D_\mu \varphi^\dagger D^\mu \varphi \\ & + \mu^2 \varphi^\dagger \varphi - \frac{\lambda}{2} (\varphi^\dagger \varphi)^2 \\ & - \bar{q}_L Y^u \tilde{\varphi} u_R - \bar{q}_L Y^d \varphi d_R - \bar{l}_L Y^e \varphi e_R + \text{H.c.} \\ & - \frac{\epsilon^{\mu\nu\alpha\beta}}{64\pi^2} \left(g_s^2 \theta\, G^a_{\mu\nu} G^a_{\alpha\beta} + g^2 \theta_w\, W^i_{\mu\nu} W^i_{\alpha\beta} + g'^2 \theta_b\, B_{\mu\nu} B_{\alpha\beta} \right), \end{aligned} \tag{6}$$

where $\epsilon^{\mu\nu\alpha\beta}$ is the totally antisymmetric symbol in four dimensions ($\epsilon^{0123} = 1$).

The first line of Eq. (6) contains the kinetic terms and self-interactions of the $SU_c(3)$, $SU_L(2)$, and $U_Y(1)$ gauge bosons. The second line contains the kinetic energy and the gauge couplings of fermions and scalars. These couplings are completely determined by gauge invariance. The terms of the third line form the scalar potential. With the parameter $\lambda > 0$ for stability and the parameter $\mu^2 > 0$, the scalar field acquires a vacuum expectation value $v = \sqrt{\mu^2/\lambda}$; the Higgs boson $h(x)$ represents fluctuations around this vacuum,

$$\varphi = \frac{v}{\sqrt{2}} U(x) \begin{pmatrix} 0 \\ 1 + \frac{h(x)}{v} \end{pmatrix}, \tag{7}$$

where $U(x)$ is an $SU(2)$ matrix, which encodes the three Goldstone bosons. The Goldstone bosons are not physical degrees of freedom, so that with a particular choice of gauge, the unitarity gauge, $U(x)$ can be set to one. In this gauge, the Goldstone bosons are "eaten" by the longitudinal polarizations of the massive vector bosons. We will often use this gauge to discuss the structure of dimension-six operators. After electroweak symmetry breaking, the scalar field kinetic energy provides a mass term for the weak gauge bosons. It is convenient to express the fields W^3_μ and B_μ in terms of the physical photon and Z-boson fields,

$$W^3_\mu = \cos\theta_W Z_\mu + \sin\theta_W A_\mu, \tag{8}$$

$$B_\mu = \cos\theta_W A_\mu - \sin\theta_W Z_\mu, \tag{9}$$

where the weak mixing angle θ_W is given, together with the proton charge $e > 0$, by the couplings g and g' via

$$g = -\frac{e}{\sin\theta_W}, \qquad g' = -\frac{e}{\cos\theta_W}. \tag{10}$$

The charged W-boson fields are defined as

$$W^\pm_\mu = \frac{1}{\sqrt{2}} \left(W^1_\mu \mp i W^2_\mu \right). \tag{11}$$

The next dimension-four operators one can write are the Yukawa couplings of the fermions to the scalar boson in the fourth line of Eq. (6), via matrices $Y^{u,d,e}$. After electroweak symmetry breaking, they generate the quark and lepton masses. By means of unitary transformations on the quark and lepton fields, it is always possible to make the fermion mass matrices diagonal and real, up to a common phase. For leptons, these transformations do not leave any trace, while for quarks the price to pay for the diagonalization of the mass matrix is that the interaction of the charged W bosons with the quarks is no longer flavor diagonal, an effect that can be obtained by replacing d^r_L in Eq. (6) by $V_{rs} d^s_L$, where V_{rs} is the unitary Cabibbo-Kobayashi-Maskawa (CKM) matrix. For three generations of quarks, the CKM matrix has one complex phase [1], which is responsible for the observed CP violation in the kaon and B-meson systems. However, the contribution of the CKM phase to nuclear EDMs is orders of magnitude smaller than the current experimental sensitivity, and we will neglect it in the rest of the paper. The second $\not{P}\not{T}$ parameter in the SM Lagrangian is the global phase of the quark mass matrices. It can be eliminated [44] by an axial rotation of all the quark fields $q_L \to e^{i\rho} q_L$, $q_R \to e^{-i\rho} q_R$. Such transformation is anomalous [45], and its net effect is to shift the coefficients of the P- and T-odd gluon operator in the fifth line of Eq. (6) from θ to $\bar\theta = \theta + n_f \rho$, where n_f is the number of quark flavors. Operators in that line are total derivatives, but, for non-Abelian gauge fields, they contribute to the action through extended field configurations, instantons [3]. The contribution of instantons is proportional to $\exp(-8\pi^2/g_i^2)$. For QCD instantons, the coupling constant g_s is large at low energies, and the instanton contribution to the action is not negligible. On the other hand, the electroweak coupling g is small, and electroweak instantons are extremely suppressed, negligible for all practical purposes. We will neglect the electroweak theta term.

The SM terms related to *CP* violation which we will be concerned with below can be summarized by

$$\mathcal{L}_4 = -\bar{\theta}\frac{g_s^2}{64\pi^2}\epsilon^{\mu\nu\alpha\beta}G^a_{\mu\nu}G^a_{\alpha\beta} - \left(1+\frac{h}{v}\right)\bar{q}_L\left(M_0 + M_3\tau_3\right)q_R + \text{H.c.}, \tag{12}$$

where $M_{0,3}$ are real, diagonal quark mass matrices.

There are good experimental and theoretical reasons to believe that the SM is an EFT, at the electroweak scale M_W, of a more fundamental theory, and it is renormalizable in the more general sense of including all higher-dimensional operators allowed by the symmetries. In the form of Eq. (6) the SM does not include neutrino masses and mixings. These can be accounted for by introducing the only gauge-invariant dimension-five operator that can be written in terms of SM fields [13]. The lightness of neutrinos compared to other fermions is explained if the new-physics scale that suppresses this operator is of the order of 10^{15} GeV. Any new physics between the electroweak and the GUT or Planck scales will manifest itself in higher-dimensional operators. A promising strategy to probe such operators is to look for processes where the SM contribution is extremely small. Examples are rare processes like the lepton flavor-changing process $\mu \to e\gamma$ or neutrinoless double beta decay.

Our work focuses on EDMs, which signal *CP* violation in the flavor-diagonal sector and are insensitive to the phase of the CKM matrix. Since $\bar{\theta}$ is very small, it is possible that higher-dimensional $\not{P}\not{T}$ operators are competitive with the $\bar{\theta}$ term. These higher-dimensional operators might eventually be linked to an underlying, ultraviolet complete $\not{P}\not{T}$ theory. We denote the scale characteristic of this theory by $M_{\not{T}}$. Well below the scale $M_{\not{T}}$ we expect $\not{P}\not{T}$ effects to be captured by the lowest-dimensional interactions among SM fields that respect the theory's gauge symmetry. In general, operators of dimension $(4+n)$ at the SM scale are suppressed by powers of $M_{\not{T}}^{-n}$. The next-to-lowest-dimension $\not{P}\not{T}$ operators involving quark and gluon fields that can be added to the SM Lagrangian have effective dimension six. Their complete set was constructed in Refs. [14, 15, 20, 16] and recently reviewed in Ref. [17].

We focus here on flavor-diagonal $\not{P}\not{T}$ operators involving quarks and gauge bosons only. Following Ref. [17], we organize the dimension-six operators we need at the electroweak scale according to their field content: three vector bosons (X^3), two gauge bosons and two scalars ($X^2\varphi^2$), two quarks, a scalar, and a vector boson ($q^2\varphi X$), two quarks, two scalars, and a derivative ($q^2\varphi^2 D$), two quarks and three scalars ($q^2\varphi^3$), and four quarks (q^4). The coefficients of these operators are all proportional to $M_{\not{T}}^{-2}$.

At dimension six, the interactions of a quark with gauge bosons gain in complexity. With the aid of the scalar boson we can write

$$\begin{aligned}\mathcal{L}_{q^2\varphi X} \;=\; & -\frac{1}{\sqrt{2}}\bar{q}_L\sigma^{\mu\nu}\left(\tilde{\Gamma}^u\lambda^a G^a_{\mu\nu} + \Gamma^u_B B_{\mu\nu} + \Gamma^u_W\tau^i W^i_{\mu\nu}\right)\frac{\tilde{\varphi}}{v}u_R \\ & -\frac{1}{\sqrt{2}}\bar{q}_L\sigma^{\mu\nu}\left(\tilde{\Gamma}^d\lambda^a G^a_{\mu\nu} + \Gamma^d_B B_{\mu\nu} + \Gamma^d_W\tau^i W^i_{\mu\nu}\right)\frac{\varphi}{v}d_R + \text{H.c.}\,,\end{aligned} \tag{13}$$

where, in the most general case, the couplings $\tilde{\Gamma}^{u,d}$ and $\Gamma^{u,d}_{B,W}$, are 3×3 complex-valued matrices. After electroweak symmetry breaking, the operators in Eq. (13) can be expressed in terms of dipole moment operators,

$$
\begin{aligned}
\mathcal{L}_{q^2\varphi X} \;=\;& -\frac{1}{2}\bar{q}_L\sigma^{\mu\nu}\left(\tilde{\Gamma}_0+\tilde{\Gamma}_3\tau_3\right)\lambda^a q_R\, G^a_{\mu\nu}-\frac{1}{2}\bar{q}_L\sigma^{\mu\nu}\left(\Gamma_0+\Gamma_3\tau_3\right)q_R\, F_{\mu\nu} \\
& -\frac{1}{2}\bar{q}_L\sigma^{\mu\nu}\left(\Gamma_{Z\,0}+\Gamma_{Z\,3}\tau_3\right)q_R\, Z_{\mu\nu} \\
& -\frac{1}{\sqrt{2}}\bar{d}_L\sigma^{\mu\nu}\Gamma^u_W u_R\, W^-_{\mu\nu}-\frac{1}{\sqrt{2}}\bar{u}_L\sigma^{\mu\nu}\Gamma^d_W d_R\, W^+_{\mu\nu}+\text{H.c.}+\ldots, \qquad (14)
\end{aligned}
$$

where, at a renormalization scale $\mu \simeq M_W$,

$$
\begin{aligned}
\tilde{\Gamma}_{0,3} &= \frac{1}{2}\left(\tilde{\Gamma}^u \pm \tilde{\Gamma}^d\right), & (15)\\
\Gamma_{0,3} &= \frac{1}{2}\left[\left(\Gamma^u_B \pm \Gamma^d_B\right)\cos\theta_W+\left(\Gamma^u_W \mp \Gamma^d_W\right)\sin\theta_W\right], & (16)\\
\Gamma_{Z\,0,3} &= \frac{1}{2}\left[\left(\Gamma^u_W \mp \Gamma^d_W\right)\cos\theta_W-\left(\Gamma^u_B \pm \Gamma^d_B\right)\sin\theta_W\right]. & (17)
\end{aligned}
$$

The first two operators in Eq. (14) are the most interesting for low-energy applications. The imaginary parts of the diagonal entries of $\Gamma_{0,3}$ and $\tilde{\Gamma}_{0,3}$ generate the quark electric and chromo-electric dipole moments (qEDM and qCEDM). Since these operators flip the chirality of the quark field, we assume these matrices to be proportional to the Yukawa couplings in the SM Lagrangian, and thus to the masses:

$$
\tilde{\Gamma}_{0,3}=\mathcal{O}\left(4\pi\tilde{\delta}_{0,3}\frac{M_0}{M^2_{\not{T}}}\right), \qquad \Gamma_{0,3}=\mathcal{O}\left(e\delta_{0,3}\frac{M_0}{M^2_{\not{T}}}\right), \qquad (18)
$$

where $\tilde{\delta}_{0,3}$ and $\delta_{0,3}$ are dimensionless constants that parameterize any deviation from this assumption, and contain all the information on physics beyond the SM. The non-diagonal entries are also of considerable interest, since they produce flavor-changing neutral currents. In the *CP*-odd sector, for example, the *uc* entries were found [46] to be the less constrained dimension-six operators that contribute to the recently observed *CP* violation in charm decays [47]. Since the flavor-changing operators contribute to nuclear EDMs only via additional loops involving weak-boson exchange, for our current purposes we can neglect them. The next three operators in Eq. (14) are weak dipole moments. Their contribution to the qEDM and qCEDM is suppressed by $(g/4\pi)^2$. In the "…" in Eq. (14) we find couplings of the Higgs boson, which we can also neglect.

The self-interaction of gauge bosons also exhibits new structures at dimension six. *CP* violation can be found in

$$
\mathcal{L}_{X^3}=\frac{d_W}{6}f^{abc}\epsilon^{\mu\nu\alpha\beta}G^a_{\alpha\beta}G^b_{\mu\rho}G^{c\,\rho}_{\nu}+\frac{d_w}{6}\epsilon^{ijk}\epsilon^{\mu\nu\alpha\beta}W^i_{\alpha\beta}W^j_{\mu\rho}W^{k\,\rho}_{\nu}. \qquad (19)
$$

The first operator is the Weinberg three-gluon operator [20] which can be interpreted as the gluon chromo-electric dipole moment (gCEDM) [21]. Similarly, the second operator is the W-boson weak electric dipole moment. After electroweak symmetry breaking, this operator generates interactions containing at least two heavy gauge bosons [15], which can contribute to q(C)EDMs through loop corrections. Again such contributions are suppressed by $(g/4\pi)^2$ [15]. For our purposes we can neglect the weak dipole moment and focus on the gCEDM,

$$
d_W=\mathcal{O}\left(4\pi\frac{w}{M^2_{\not{T}}}\right), \qquad (20)
$$

with w a dimensionless constant.

Just as there is CP violation in multi-gluon operators, so there is in multi-quark operators. In agreement with Ref. [16], we find just two four-quark interactions,

$$\mathcal{L}_{q^4} = \Sigma_1 \left(\bar{q}_L^J u_R\right) \epsilon_{JK} \left(\bar{q}_L^K d_R\right) + \Sigma_8 \left(\bar{q}_L^J \lambda^a u_R\right) \epsilon_{JK} \left(\bar{q}_L^K \lambda^a d_R\right) + \text{H.c.}, \tag{21}$$

where the couplings $\Sigma_{1,8}$ are four-index tensors in flavor space. They scale as

$$\Sigma_{1,8} = \mathcal{O}\left((4\pi)^2 \frac{\sigma_{1,8}}{M_{\not T}^2}\right), \tag{22}$$

where $\sigma_{1,8}$ are dimensionless constants. The operators in Eq. (21) are not affected by electroweak symmetry breaking. For later convenience, we rewrite Eq. (21) in terms of quark doublets $q = q_L + q_R$, and focus on the flavor-diagonal $\not P \not T$ terms only,

$$\begin{aligned} \mathcal{L}_{q^4} \;\; = \;\; & \frac{1}{4} \left(\text{Im}\Sigma_1\right)_{1111} \left(\bar{q}q\, \bar{q}i\gamma_5 q - \bar{q}\boldsymbol{\tau}q \,\cdot \bar{q}\boldsymbol{\tau}i\gamma_5 q\right) \\ & + \frac{1}{4} \left(\text{Im}\Sigma_8\right)_{1111} \left(\bar{q}\lambda^a q\, \bar{q}i\gamma_5\lambda^a q - \bar{q}\boldsymbol{\tau}\lambda^a q \,\cdot \bar{q}\boldsymbol{\tau}i\gamma_5\lambda^a q\right) + \dots \end{aligned} \tag{23}$$

These four-quark operators, originating directly at the electroweak scale and constrained by exact $SU_L(2)$ gauge invariance, are chiral invariant. The remaining parts of Eq. (21) and other dimension-six $\not P \not T$ four-quark operators listed in Ref. [17] necessarily involve flavor-changing effects, and contribute to flavor-diagonal CP violation only at the loop level. We will neglect them in what follows.

The remaining $\not P \not T$ sources involve two or more scalar fields. The most important of them is

$$\mathcal{L}_{q^2\varphi^2 D} = \bar{u}_R \Xi_1 \gamma^\mu d_R \; \tilde{\varphi}^\dagger i D_\mu \varphi + \text{H.c.}, \tag{24}$$

where in general Ξ_1 is a complex 3×3 matrix in flavor space. Here we focus, again, on the flavor-diagonal parts only, given after electroweak symmetry breaking by

$$\mathcal{L}_{q^2\varphi^2 D} = \frac{g v^2}{2\sqrt{2}} \left(W_\mu^+ \bar{u}_R \Xi_1 \gamma^\mu d_R + W_\mu^- \bar{d}_R \Xi_1 \gamma^\mu u_R\right) + \dots \tag{25}$$

Contrary to the operators in Eq. (13), the operators in Eq. (25) do not change chirality and we do not expect them to be proportional to the quark mass, so we parameterize

$$\Xi_1 = \mathcal{O}\left((4\pi)^2 \frac{\xi}{M_{\not T}^2}\right). \tag{26}$$

At low energy, after we integrate out the W boson, the imaginary part of Ξ_1 contributes to $\not P \not T$ four-quark operators, which are particularly interesting because they are not suppressed by the light quark masses [22]. As detailed in Sec. 3, $\text{Im}\,\Xi_1$ generates four-quark operators of the same importance as the chiral-invariant four-quark operators that are generated directly at the electroweak scale, which we introduced in Eq. (21).

Finally, there are other terms that are closely related to interactions in Eq. (6),

$$\begin{aligned} \mathcal{L}_{X^2\varphi^2, q^2\varphi^3} \;\; = \;\; & -\frac{\epsilon^{\mu\nu\alpha\beta}}{32\pi^2} \left(g_s^2 \theta'\, G^a_{\mu\nu} G^a_{\alpha\beta} + g^2 \theta'_w\, W^i_{\mu\nu} W^i_{\alpha\beta} + g'^2 \theta'_b\, B_{\mu\nu} B_{\alpha\beta}\right) \frac{\varphi^\dagger \varphi}{v^2} \\ & + \frac{\epsilon^{\mu\nu\alpha\beta}}{32\pi^2} g g' \theta'_{wb}\, W^i_{\mu\nu} B_{\alpha\beta} \frac{\varphi^\dagger \tau^i \varphi}{v^2} - 2 \frac{\varphi^\dagger \varphi}{v^2} \left(\bar{q}_L Y'^u \tilde{\varphi} u_R + \bar{q}_L Y'^d \tilde{\varphi} d_R\right), \end{aligned} \tag{27}$$

where the angles θ', $\theta'_{w,\,b,\,wb}$ and the Yukawa couplings $Y'^{\,u,d}$, which are symmetric, complex matrices in flavor space, scale as

$$\theta', \theta'_{w,\,b,\,wb} = \mathcal{O}\left(\frac{v^2}{M_{\not T}^2}\right), \qquad Y'^{\,u,d} = \mathcal{O}\left(\frac{v^2}{M_{\not T}^2}\right). \tag{28}$$

Equation (27) is reminiscent of Eq. (6). Indeed, if one rewrites $2\varphi^\dagger\varphi/v^2 = 1 + (2\varphi^\dagger\varphi/v^2 - 1)$, the pieces of the first three and last two operators that do not contain the Higgs boson can be absorbed in a redefinition of the couplings θ, $\theta_{w,\,b}$, and $Y^{u,d}$ in Eq. (6). In some sense this aggravates the strong *CP* problem since even if the bare QCD vacuum angle is tuned to zero, we need to explain why the contributions induced by these higher-dimensional terms are small as well. Of the first four terms in Eq. (27), only the operator θ'_{wb} obtains after electroweak breaking a nontopological piece without a Higgs boson [15]. This piece connects three or more electroweak gauge bosons and contributes through loop diagrams to the quark electric and weak dipole moments, but such contributions are suppressed by $(g/4\pi)^2$. The remainder of Eq. (27) then includes *CP*-odd operators with at least one Higgs boson: in the unitarity gauge,

$$\begin{aligned}\mathcal{L}_{X^2\varphi^2,q^2\varphi^3} \;=\; -\Bigg\{ & \frac{\epsilon^{\mu\nu\alpha\beta}}{32\pi^2}\left(g_s^2\theta'\, G^a_{\mu\nu}G^a_{\alpha\beta} + g^2\theta'_w\, W^i_{\mu\nu}W^i_{\alpha\beta} + g'^2\theta'_b\, B_{\mu\nu}B_{\alpha\beta} + gg'\theta'_{wb}\, W^3_{\mu\nu}B_{\alpha\beta}\right) \\ & + \sqrt{2}v\left(1+\frac{h}{v}\right)\left(\bar u_L Y'^{\,u} u_R + \bar d_L Y'^{\,d} d_R\right)\Bigg\}\frac{h}{v}\left(1+\frac{h}{2v}\right).\end{aligned} \tag{29}$$

In the first line of Eq. (29) are *CP*-odd interactions of the Higgs to two gluons, two photons, or two weak bosons. The imaginary parts of the Yukawa couplings generate flavor-diagonal and flavor-changing $\not P \not T$ Higgs-quark interactions. At low energy, where the Higgs boson is integrated out, these interactions manifest themselves in loop corrections to the qEDM, qCEDM and gCEDM. At tree level, $\not P \not T$ Higgs-quark interactions generate $\not P \not T$ four-quark operators, which, as discussed below, are relatively small.

Of course, we could continue the procedure to higher-dimensional $\not P \not T$ sources. Those would include, for example, further four-quark operators [48] at dimension eight. We limit ourselves here to a systematic study of the effects of dimension-six sources to low-energy observables.

3 Tree-level matching onto the QCD scale

The $\not P \not T$ Lagrangian of dimension up to six that is relevant for the calculation of hadronic and nuclear EDMs is summarized in Eqs. (12), (14), (19), (23), (25), and (29). For low-energy applications, it is important to evolve the $\not P \not T$ Lagrangian from the electroweak scale down to the typical hadronic scale $\mu \sim M_{\rm QCD} \simeq 1$ GeV. In the process, one has to integrate out the effects of heavy SM particles [15, 49]; at the same time, one has to evaluate the running of the coupling constants and account for the possible mixing of the dimension-six operators [21, 50, 48]. A detailed account of the matching and evolution of the complete dimension-six $\not P \not T$ Lagrangian is beyond the scope of this work. Here we limit ourselves to tree-level matching, and we turn off (most of) the running of the couplings.

At the QCD scale, the theory involves only light quarks, gluons and photons. Focusing on the first generation,

$$\mathcal{L}_{\rm cl} = -\frac{1}{4}\left(G^a_{\mu\nu}G^{a\,\mu\nu} + F_{\mu\nu}F^{\mu\nu}\right) + \bar q\left(i\not\partial + \frac{g_s}{2}\not G^a\lambda^a\right)q \tag{30}$$

has a global, chiral $SU_L(2) \times SU_R(2) \sim SO(4)$ symmetry under independent $SU(2)$ rotations of left- and right-handed quarks. Remaining dimension-four (and most of the higher-dimensional) interactions break chiral symmetry explicitly. Because all the breaking is due to relatively small quantities, it is useful to classify interactions according to their transformation pattern under $SO(4)$.

For the construction of $\not{P}\not{T}$ electromagnetic operators in the chiral Lagrangian, we will need to consider the chiral properties of the PT electromagnetic couplings of the quarks in

$$\mathcal{L}_e = -eA_\mu \bar{q}\gamma^\mu Q q, \tag{31}$$

where

$$Q = \frac{1}{6} + \frac{\tau_3}{2} \tag{32}$$

is the quark charge matrix. Introducing a chiral-invariant I^μ and an antisymmetric tensor T^μ through

$$I^\mu = \bar{q}\gamma^\mu q, \qquad T^\mu = \begin{pmatrix} \epsilon^{ijk}\bar{q}\gamma^\mu\gamma_5\tau^k q & \bar{q}\gamma^\mu\tau^j q \\ -\bar{q}\gamma^\mu\tau^i q & 0 \end{pmatrix}, \tag{33}$$

we rewrite Eq. (31) as

$$\mathcal{L}_e = -\frac{e}{2}A_\mu \left(\frac{I^\mu}{3} + T^\mu_{34}\right). \tag{34}$$

At low energy, we can neglect the effects of the Higgs and of the heavy quarks, and Eq. (12) is rewritten as

$$\mathcal{L}_4 = -\bar{q}_L M q_R - \bar{q}_R M q_L - \bar{\theta}\frac{g_s^2}{64\pi^2}\epsilon^{\mu\nu\alpha\beta}\, G^a_{\mu\nu}G^a_{\alpha\beta}\ , \tag{35}$$

where

$$M = \begin{pmatrix} m_u & 0 \\ 0 & m_d \end{pmatrix} = \bar{m}\left(1 - \varepsilon\tau_3\right), \tag{36}$$

with real parameters $m_{u,d}$, or alternatively

$$\bar{m} = \frac{m_u + m_d}{2}, \qquad \varepsilon = \frac{m_d - m_u}{m_u + m_d}. \tag{37}$$

For a χPT treatment, it is more convenient to eliminate the $\bar{\theta}$ term with an axial $U(1)$ rotation on the quark fields, and move all CP violation to the quark mass term. After vacuum alignment [44], the QCD $\bar{\theta}$ term becomes, in the notation of Ref. [32],

$$\mathcal{L}_4 = \bar{m}r(\bar{\theta})S_4 - \varepsilon\bar{m}\, r^{-1}(\bar{\theta})P_3 - m_*\sin\bar{\theta}\, r^{-1}(\bar{\theta})P_4\ , \tag{38}$$

where we introduced two $SO(4)$ vectors

$$S = \begin{pmatrix} -i\bar{q}\gamma^5\boldsymbol{\tau}q \\ \bar{q}q \end{pmatrix}, \qquad P = \begin{pmatrix} \bar{q}\boldsymbol{\tau}q \\ i\bar{q}\gamma^5 q \end{pmatrix}, \tag{39}$$

a function $r(\bar{\theta})$ that goes to 1 in the limit of small $\bar{\theta}$,

$$r(\bar{\theta}) = \left(\frac{1 + \varepsilon^2\tan^2\frac{1}{2}\bar{\theta}}{1 + \tan^2\frac{1}{2}\bar{\theta}}\right)^{1/2} \simeq 1 + \mathcal{O}(\bar{\theta}^2), \tag{40}$$

and the parameter

$$m_* = \frac{m_u m_d}{m_u + m_d} = \frac{1-\varepsilon^2}{2}\bar{m}. \tag{41}$$

From these ingredients one can derive the form of low-energy interactions originating in the QCD vacuum angle, as presented, for example, in Ref. [32].

We will now consider the dimension-six Lagrangian, which breaks chiral symmetry in new ways. The light-flavor components of the dipole operators in Eq. (14) match onto the light-quark EDM and CEDM,

$$\mathcal{L}_{\rm q(C)EDM} = -\frac{1}{2}\,\bar{q}\, i\sigma^{\mu\nu}\gamma^5\left(d_0 + d_3\tau_3\right) q\, F_{\mu\nu} - \frac{1}{2}\,\bar{q}\, i\sigma^{\mu\nu}\gamma^5\left(\tilde{d}_0 + \tilde{d}_3\tau_3\right)\lambda^a q\, G^a_{\mu\nu}, \tag{42}$$

where, at tree level,

$$d_{0,3} = {\rm Im}(\Gamma_{0,3})_{11}, \qquad \tilde{d}_{0,3} = {\rm Im}(\tilde{\Gamma}_{0,3})_{11}. \tag{43}$$

The non-diagonal components of the qEDM and qCEDM operators in Eq. (14) correct Eq. (43) at the loop level. The weak EDMs also generate loop corrections to the qEDM and qCEDM. At tree level, they generate four-quark operators, which, however, involve one power of the quark momentum, and are thus dimension-seven. The couplings of such operators are suppressed by an additional factor $m_{u,d}/M_W^2$ compared to the couplings of the dimension-six operators that we keep.

The gCEDM in Eq. (19) and the four-quark operators in Eq. (23) match onto themselves. As pointed out in Ref. [22], at tree level, the only non-trivial result comes from the operator in Eq. (25). The largest effect comes from a W exchange between light quarks, the suppression by M_W^2 from the W propagator being compensated by the factor g^2v^2 from the vertices. The resulting $\not{P}\not{T}$ four-quark operator, when evolved from M_W to $M_{\rm QCD}$, induces another four-quark operator of similar form but with additional color structure [1]. The latter does not appear directly at the electroweak scale due to its gauge-symmetry-breaking properties. We find the following $\not{P}\not{T}$ operators, which because of their left-right mixing we abbreviate as FQLR:

$$\begin{aligned}\mathcal{L}_{\rm LR} \;=\;& i\,{\rm Im}(\Xi_1)_{11}\, V_{ud}\left(\bar{u}_R\gamma^\mu d_R\, \bar{d}_L\gamma_\mu u_L - \bar{d}_R\gamma^\mu u_R\, \bar{u}_L\gamma_\mu d_L\right) \\ &+ i\,{\rm Im}(\Xi_8)_{11}\, V_{ud}\left(\bar{u}_R\gamma^\mu\lambda^a d_R\, \bar{d}_L\gamma_\mu\lambda^a u_L - \bar{d}_R\gamma^\mu\lambda^a u_R\, \bar{u}_L\gamma_\mu\lambda^a d_L\right),\end{aligned} \tag{44}$$

where $V_{ud} \simeq 1$ is a CKM element. Here, $(\Xi_8)_{11}$ is not an independent coupling, but it depends on $(\Xi_1)_{11}$ and on QCD renormalization-group factors, which are calculable. Thus $(\Xi_1)_{11}$ and $(\Xi_8)_{11}$ both depend on the same dimensionless parameter ξ. In terms of light-quark doublets, we can rewrite Eq. (44) as

$$\mathcal{L}_{\rm LR} = \frac{1}{4}\,{\rm Im}(\Xi_1)_{11}\,\epsilon^{3ij}\,\bar{q}\tau^i\gamma^\mu q\,\bar{q}\tau^j\gamma_\mu\gamma_5 q + \frac{1}{4}\,{\rm Im}(\Xi_8)_{11}\,\epsilon^{3ij}\,\bar{q}\tau^i\gamma^\mu\lambda^a q\,\bar{q}\tau^j\gamma_\mu\gamma_5\lambda^a q. \tag{45}$$

By a Fierz rearrangement we can further rewrite the first term of this equation as

$$\frac{1}{8}\,{\rm Im}(\Xi_1)_{11}\left[2\left(\bar{q}q\,\bar{q}i\gamma_5\tau_3 q - \bar{q}\tau_3 q\,\bar{q}i\gamma_5 q\right) - 3\left(\bar{q}\lambda^a q\,\bar{q}i\gamma_5\tau_3\lambda^a q - \bar{q}\tau_3\lambda^a q\,\bar{q}i\gamma_5\lambda^a q\right)\right]. \tag{46}$$

This operator was studied in Ref. [51] in the framework of left-right models. Although a large list of four-quark operators is presented there, only one combination ($O_{11} - O_{12} + 6O_{21} - 6O_{22}$, in their notation) is generated at the electroweak scale. This combination is identical to Eq. (44). It is interesting to point out that we could have coupled the operator in Eq. (25) to the left-handed lepton current. The operator created this way causes $\not{T}$ in β decay through contributions to the triple correlation $\sim D\,\vec{J}\cdot(\vec{p}_e \times \vec{p}_\nu)$. In Ref. [22] it is argued that, with current experimental accuracy, the best limit on the FQLR operator comes from EDM experiments.

---

[1] This was pointed out to us by W. Dekens. We thank him for useful discussions on this subject.

All other $\not{P}\not{T}$ effects have extra suppression at low energies. The operators in Eq. (29) contain at least one Higgs field, and generate $\not{P}\not{T}$ effects at either tree or loop level. For example, at tree level, the operators in the second line contribute to four-quark operators. But, since the Higgs couples to the light quark mass, the resulting $\not{P}\not{T}$ operators have a suppression of at least the small ratio of the light-quark mass over the electroweak scale.

In summary, the (effectively) dimension-six $\not{P}\not{T}$ Lagrangian is

$$\begin{aligned}
\mathcal{L}_6 \;=\;& -\frac{1}{2}\bar{q}\,(d_0 + d_3\tau_3)\, i\sigma^{\mu\nu}\gamma_5 q\, F_{\mu\nu} - \frac{1}{2}\bar{q}\left(\tilde{d}_0 + \tilde{d}_3\tau_3\right) i\sigma^{\mu\nu}\gamma_5\lambda^a q\, G^a_{\mu\nu} \\
& + \frac{d_W}{6} f^{abc}\epsilon^{\mu\nu\alpha\beta} G^a_{\alpha\beta} G^b_{\mu\rho} G^{c\,\rho}_{\nu} \\
& + \frac{\mathrm{Im}\,\Xi_1}{4}\epsilon^{3ij}\,\bar{q}\tau^i\gamma^\mu q\,\bar{q}\tau^j\gamma_\mu\gamma_5 q + \frac{\mathrm{Im}\,\Xi_8}{4}\epsilon^{3ij}\,\bar{q}\tau^i\gamma^\mu\lambda^a q\,\bar{q}\tau^j\gamma_\mu\lambda^a\gamma_5 q \\
& + \frac{\mathrm{Im}\Sigma_1}{4}\left(\bar{q}q\,\bar{q}i\gamma_5 q - \bar{q}\boldsymbol{\tau} q\,\cdot\bar{q}\boldsymbol{\tau} i\gamma_5 q\right) + \frac{\mathrm{Im}\Sigma_8}{4}\left(\bar{q}\lambda^a q\,\bar{q}i\gamma_5\lambda^a q - \bar{q}\boldsymbol{\tau}\lambda^a q\,\cdot\bar{q}\boldsymbol{\tau} i\gamma_5\lambda^a q\right), \qquad (47)
\end{aligned}$$

where all coupling constants have been redefined in order to absorb effects from operator mixing and renormalization-group running, and to drop the generation indices in the case of four-quark operators. As discussed above, one can imagine additional $\not{P}\not{T}$ four-quark operators, but such operators are suppressed by weak gauge couplings in the typical combination $(g/4\pi)^2$, small off-diagonal CKM elements, and/or powers of $m_{u,d}/M_W$. We therefore expect the operators in Eqs. (38) and (47) to give rise to the dominant P and T violation in hadronic and nuclear systems at low energy.

The redefined constants in Eq. (47) scale as

$$\begin{aligned}
& d_{0,3} = \mathcal{O}\left(\frac{e\delta_{0,3}\bar{m}}{M^2_{\not{T}}}\right), \qquad \tilde{d}_{0,3} = \mathcal{O}\left(4\pi\frac{\tilde{\delta}_{0,3}\bar{m}}{M^2_{\not{T}}}\right), \qquad d_W = \mathcal{O}\left(4\pi\frac{w}{M^2_{\not{T}}}\right), \\
& \Xi_{1,8} = \mathcal{O}\left(\frac{(4\pi)^2\xi}{M^2_{\not{T}}}\right), \qquad \Sigma_{1,8} = \mathcal{O}\left(\frac{(4\pi)^2\sigma_{1,8}}{M^2_{\not{T}}}\right), \qquad (48)
\end{aligned}$$

in terms of dimensionless numbers $\delta_{0,3}$, $\tilde{\delta}_{0,3}$, w, ξ, and $\sigma_{1,8}$. The sizes of these dimensionless parameters depend on the exact mechanisms of electroweak and P and T breaking, and on the running to the low energies where non-perturbative QCD effects take over. The minimal assumption is that $\delta_{0,3}$, $\tilde{\delta}_{0,3}$, w, ξ, and $\sigma_{1,8}$ are $\mathcal{O}(1)$, $\mathcal{O}(g_s/4\pi)$, $\mathcal{O}((g_s/4\pi)^3)$, $\mathcal{O}(1)$, and $\mathcal{O}(1)$, respectively. However, they can be much smaller or larger, depending on the parameters encoding $\not{P}\not{T}$ beyond the Standard Model. In the SM itself, where $M_{\not{T}} = M_W$, $\delta_{0,3}$, $\tilde{\delta}_{0,3}$ and w are suppressed not only by the Jarlskog parameter [2] $J_{\mathrm{CP}} \simeq 3\times 10^{-5}$, but also by additional powers of small gauge coupling constants and ratios of quark-to-W masses [52, 7]. In certain supersymmetric models with various simplifying universality assumptions of a soft-breaking sector with a common scale M_{SUSY}, one has $M_{\not{T}} = M_{\mathrm{SUSY}}$ and the dimensionless parameters are of the size of the minimal assumption times a factor which is [7, 53], roughly, $A_{CP} = (g_s/4\pi)^2\sin\phi$ (neglecting electroweak parameters), with ϕ a phase encoding T violation. If in the soft-breaking sfermion mass matrices non-diagonal terms are allowed, enhancements of the type $m_b/m_d \sim 10^3$ or even $m_t/m_u \sim 10^5$ become possible, although they are usually associated with other, smaller phases [7]. In other models, the relative sizes of the dimensionless parameters could be different still.

Below the hadronic scale $M_{\rm QCD}$, the dimension-six $\not{P}\not{T}$ sources generate further effective interactions, which break chiral symmetry in their own ways. Introducing the $SO(4)$ singlets

$$\begin{aligned} I_W &= \frac{1}{6} f^{abc}\epsilon^{\mu\nu\alpha\beta} G^a_{\alpha\beta} G^b_{\mu\rho} G^{c\,\rho}_{\nu}, \\ I^{(1)}_{qq} &= \frac{1}{4}\left(\bar{q}q\,\bar{q}i\gamma_5 q - \bar{q}\boldsymbol{\tau}q\cdot\bar{q}\boldsymbol{\tau}i\gamma_5 q\right) = \frac{1}{4} S\cdot P, \\ I^{(8)}_{qq} &= \frac{1}{4}\left(\bar{q}\lambda^a q\,\bar{q}i\gamma_5\lambda^a q - \bar{q}\boldsymbol{\tau}\lambda^a q\cdot\bar{q}\boldsymbol{\tau}i\gamma_5\lambda^a q\right), \end{aligned} \tag{49}$$

the $SO(4)$ vectors

$$W = \tfrac{1}{2}\begin{pmatrix} -i\bar{q}\sigma^{\mu\nu}\gamma_5\boldsymbol{\tau}q \\ \bar{q}\sigma^{\mu\nu}q \end{pmatrix} F_{\mu\nu}, \qquad V = \tfrac{1}{2}\begin{pmatrix} \bar{q}\sigma^{\mu\nu}\boldsymbol{\tau}q \\ i\bar{q}\sigma^{\mu\nu}\gamma_5 q \end{pmatrix} F_{\mu\nu}, \tag{50}$$

and

$$\tilde{W} = \tfrac{1}{2}\begin{pmatrix} -i\bar{q}\sigma^{\mu\nu}\gamma_5\boldsymbol{\tau}\lambda^a q \\ \bar{q}\sigma^{\mu\nu}\lambda^a q \end{pmatrix} G^a_{\mu\nu}, \qquad \tilde{V} = \tfrac{1}{2}\begin{pmatrix} \bar{q}\sigma^{\mu\nu}\boldsymbol{\tau}\lambda^a q \\ i\bar{q}\sigma^{\mu\nu}\gamma_5\lambda^a q \end{pmatrix} G^a_{\mu\nu}, \tag{51}$$

and the symmetric $SO(4)$ tensors

$$X^{(1)} = \frac{1}{4}\begin{pmatrix} \bar{q}\tau^i\gamma^\mu q\,\bar{q}\tau^j\gamma_\mu q - \bar{q}\tau^i\gamma^\mu\gamma_5 q\,\bar{q}\tau^j\gamma_\mu\gamma_5 q & -\epsilon^{jkl}\bar{q}\tau^k\gamma^\mu q\,\bar{q}\tau^l\gamma_\mu\gamma_5 q \\ -\epsilon^{ikl}\bar{q}\tau^k\gamma^\mu q\,\bar{q}\tau^l\gamma_\mu\gamma_5 q & \bar{q}\boldsymbol{\tau}\gamma^\mu q\cdot\bar{q}\boldsymbol{\tau}\gamma_\mu q - \bar{q}\boldsymbol{\tau}\gamma^\mu\gamma_5 q\cdot\bar{q}\boldsymbol{\tau}\gamma_\mu\gamma_5 q \end{pmatrix}, \tag{52}$$

and

$$X^{(8)} = \frac{1}{4}\begin{pmatrix} \bar{q}\tau^i\gamma^\mu\lambda^a q\,\bar{q}\tau^j\gamma_\mu\lambda^a q - \bar{q}\tau^i\gamma^\mu\gamma_5\lambda^a q\,\bar{q}\tau^j\gamma_\mu\gamma_5\lambda^a q & -\epsilon^{jkl}\bar{q}\tau^k\gamma^\mu\lambda^a q\,\bar{q}\tau^l\gamma_\mu\gamma_5\lambda^a q \\ -\epsilon^{ikl}\bar{q}\tau^k\gamma^\mu\lambda^a q\,\bar{q}\tau^l\gamma_\mu\gamma_5\lambda^a q & \bar{q}\boldsymbol{\tau}\gamma^\mu\lambda^a q\cdot\bar{q}\boldsymbol{\tau}\gamma_\mu\lambda^a q - \bar{q}\boldsymbol{\tau}\gamma^\mu\gamma_5\lambda^a q\cdot\bar{q}\boldsymbol{\tau}\gamma_\mu\gamma_5\lambda^a q \end{pmatrix}, \tag{53}$$

we summarize the $\not{P}\not{T}$ Lagrangian as

$$\mathcal{L}_6 = -d_0 V_4 + d_3 W_3 - \tilde{d}_0\tilde{V}_4 + \tilde{d}_3\tilde{W}_3 + d_W I_W - \mathrm{Im}\,\Xi_1\,X^{(1)}_{34} - \mathrm{Im}\,\Xi_8\,X^{(8)}_{34} + \mathrm{Im}\Sigma_1\,I^{(1)}_{qq} + \mathrm{Im}\Sigma_8\,I^{(8)}_{qq}. \tag{54}$$

We already emphasized that the dimensionless coefficients $\delta_{0,3}$, $\tilde{\delta}_{0,3}$, w, ξ, and $\sigma_{1,8}$ that we have introduced, and their relative sizes, strongly depend on the particular high-energy model, making it difficult to compare the relative contributions of different $\not{P}\not{T}$ sources to the same observable in a way that is independent of the details of the physics at the high-energy scale $M_{\not{T}}$. We therefore adopt the approach that we construct the low-energy $\not{P}\not{T}$ Lagrangian for each dimension-six source separately. These separate Lagrangians can be used to calculate for each source their contribution to hadronic $\not{P}\not{T}$ observables. Each $\not{P}\not{T}$ source generates a characteristic pattern of relations between different observables, rooted in its field content and its transformation properties under chiral symmetry. The observation of such pattern in the current generation of, for example, nucleon and nuclear EDM experiments would then effectively pinpoint the dominant $\not{P}\not{T}$ mechanism at the QCD scale [43, 33, 34, 35, 36, 37], if one exists. Once this is known, the next step would be to trace the dominant effects up to the electroweak scale.

On the other hand, the formalism we develop can be easily adjusted to specific extensions of the SM. Any low-energy observable can be obtained by combining hadronic contributions from each separate source. Once the values of $\delta_{0,3}$, $\tilde{\delta}_{0,3}$, w, ξ, and $\sigma_{1,8}$ in a given model, and their running from $M_{\not T}$ to $M_{\rm QCD}$, are known, then the relative importance of the interactions constructed in the next sections can be reassessed to accommodate, for instance, a large hierarchy between these parameters.

4 SO(4) chiral framework

Below the QCD scale quarks and gluons are no longer convenient degrees of freedom to describe strong interactions, which can be rewritten in terms of hadronic EFTs. The implications of the Lagrangian in Eqs. (30), (34), (38), and (54) for the interactions among pions and nucleons (and delta isobars, since they are not much more massive than nucleons) at low momentum $Q \sim m_\pi \ll M_{\rm QCD}$, where $m_\pi \simeq 140$ MeV is the pion mass, are described by chiral EFT, an extension to arbitrary number of nucleons of χPT. At such momenta, pions must explicitly be accounted for in the theory, while other mesons can be integrated out. At lower momenta, $Q \ll m_\pi$, even pions (and delta isobars) can be integrated out, the corresponding EFT being called pionless EFT. This EFT finds applications in loosely bound nuclei, where it is much simpler to deploy than chiral EFT for it involves only short-range inter-nucleon interactions. However, the constraints from chiral symmetry, and consequently some predictive power, are lost. The *form* of the relevant pionless EFT interactions can be read from the following by discarding pion fields.

The special role of the pion is a consequence of the approximate invariance of the QCD Lagrangian under the chiral symmetry $SU_L(2) \times SU_R(2) \sim SO(4)$. Because it is not manifest in the spectrum, which only exhibits approximate isospin symmetry, chiral symmetry must be spontaneously broken down to its diagonal, isospin subgroup, $SU_V(2) \sim SO(3)$. From Goldstone's theorem, one expects to find in the spectrum three massless Goldstone bosons that live on the "chiral circle" $S^3 \sim SO(4)/SO(3)$. The explicit breaking of chiral symmetry gives pions a mass, but the way interactions at the quark/gluon level transform constrains pion interactions. In this section we show how these constraints arise in the CP-even sector for one particular choice of pion fields, and in the next section apply this technique to $\not P \not T$ interactions. (Because observables are independent of the choice of fields, the method we use can be straightforwardly reproduced for other choices without affecting the physics.) Generalization to $SU_L(3) \times SU_R(3)$ is possible, but the $SU_L(2) \times SU_R(2)$ case is best adapted to nuclear physics, on which scale the strange quark is not particularly light. The method used in this section stems from Refs. [25, 54].

We parametrize the chiral circle with stereographic coordinates [25], whose dimensionless fields we denote by an isovector field $\boldsymbol{\zeta}$. We can identify these degrees of freedom with canonically normalized pion fields $\boldsymbol{\pi} = F_\pi \boldsymbol{\zeta}$, where $F_\pi \simeq 186$ MeV, the pion decay constant, is the diameter of the chiral circle. Such fields transform in a complicated way under chiral symmetry. However, a pion covariant derivative can be defined by

$$D_\mu \boldsymbol{\pi} = D^{-1} \partial_\mu \boldsymbol{\pi}, \tag{55}$$

with

$$D = 1 + \frac{\boldsymbol{\pi}^2}{F_\pi^2}, \tag{56}$$

which transforms under chiral transformations as under an isospin transformation, but with a field-dependent parameter. Similarly, we can use an isospin-1/2 nucleon field $N = (p\ n)^T$ that transforms in an analogous way, and a nucleon covariant derivative,

$$\mathcal{D}_\mu N = \left(\partial_\mu + \frac{i}{F_\pi^2}\boldsymbol{\tau}\cdot\boldsymbol{\pi}\times D_\mu\boldsymbol{\pi}\right)N. \tag{57}$$

We define $\mathcal{D}^\dagger$ through $\bar{N}\mathcal{D}^\dagger \equiv \overline{\mathcal{D}N}$ and use the short-hand notation:

$$\begin{aligned} \mathcal{D}^\mu_\pm &\equiv \mathcal{D}^\mu \pm \mathcal{D}^{\dagger\mu}, & \mathcal{D}^\mu_\pm\mathcal{D}^\nu_\pm &\equiv \mathcal{D}^\mu\mathcal{D}^\nu + \mathcal{D}^{\dagger\nu}\mathcal{D}^{\dagger\mu} \pm \mathcal{D}^{\dagger\mu}\mathcal{D}^\nu \pm \mathcal{D}^{\dagger\nu}\mathcal{D}^\mu, \\ \tau_i\mathcal{D}^\mu_\pm &\equiv \tau_i\mathcal{D}^\mu \pm \mathcal{D}^{\dagger\mu}\tau_i, & \tau_i\mathcal{D}^\mu_\pm\mathcal{D}^\nu_\pm &\equiv \tau_i\mathcal{D}^\mu\mathcal{D}^\nu + \mathcal{D}^{\dagger\nu}\mathcal{D}^{\dagger\mu}\tau_i \pm \mathcal{D}^{\dagger\mu}\tau_i\mathcal{D}^\nu \pm \mathcal{D}^{\dagger\nu}\tau_i\mathcal{D}^\mu. \end{aligned} \tag{58}$$

Covariant derivatives of covariant derivatives can be constructed similarly, for example

$$(\mathcal{D}_\mu D_\nu\pi)_i = \left(\partial_\mu\delta_{ij} - \frac{2}{F_\pi^2}\epsilon_{ikj}(\boldsymbol{\pi}\times D_\mu\boldsymbol{\pi})_k\right)D_\nu\pi_j. \tag{59}$$

For simplicity we omit the delta isobar here, but one can introduce an isospin-3/2 field for it along completely analogous lines [54].

Since nucleons are essentially nonrelativistic for $Q \ll m_N$, the nucleon mass, we work in the heavy-baryon framework [41] where, instead of gamma matrices, it is the nucleon velocity v^μ and spin S_μ ($S = (\vec{\sigma}/2, 0)$ in the rest frame $v = (\vec{0}, 1)$) that appear in interactions. Below we use a subscript $\perp$ to denote the component of a four-vector perpendicular to the velocity, for example

$$\mathcal{D}^\mu_\perp = \mathcal{D}^\mu - v^\mu v\cdot\mathcal{D}. \tag{60}$$

The interactions we construct are manifestly invariant under rotations and translations, but not under Lorentz boosts. Nevertheless, Lorentz invariance imposes non-trivial constraints on the interactions in the effective Lagrangian and on their coefficients [42].

The first step in describing QCD at low energies is to construct the most general Lagrangian that transforms under the symmetries of QCD in the same way as the QCD Lagrangian itself. Along with this, one needs a power-counting scheme so that interactions can be ordered according to the expected size of their contributions. The Lagrangian contains an infinite number of terms that we group using an integer "chiral index" Δ [23] and the (even) number of fermion fields f,

$$\mathcal{L} = \sum_{\Delta=0}^{\infty}\sum_{f/2}\mathcal{L}_f^{(\Delta)}. \tag{61}$$

We will restrict ourselves here mostly to $f \leq 2$, leaving a discussion of the case $f \geq 4$ for Sec. 10.

The technology for constructing the Lagrangian is well known, see, for example, Ref. [25]. When we neglect $\mathcal{L}_e$ (34), $\mathcal{L}_4$ (38), and $\mathcal{L}_6$ (54), the EFT Lagrangian includes all chiral-invariant interactions made out of $D_\mu\boldsymbol{\pi}$, N, and their covariant derivatives. In this case, $f \leq 2$ interactions have chiral index [23]

$$\Delta = d + f/2 - 2 \geq 0, \tag{62}$$

in terms of the number d of derivatives (and powers of the delta-nucleon mass difference). The coefficients of the effective operators, the so-called low-energy constants (LECs), cannot yet be calculated directly from QCD, but they can be estimated using naive dimensional analysis (NDA) [55, 20], in which case the index Δ tracks the number of inverse powers of $M_{\rm QCD} \sim 2\pi F_\pi \simeq 1.2$ GeV associated with an interaction. (Note that since NDA associates the LECs of chiral-invariant operators to $g_s/4\pi$, for consistency one should take $g_s \sim 4\pi$.) For the purposes of the present work, we need explicitly only the leading *CP*-even interactions,

$$\mathcal{L}^{(0)}_{\chi,f\leq 2} = \frac{1}{2}D_\mu \boldsymbol{\pi}\cdot D^\mu \boldsymbol{\pi} + \bar{N}\left(iv\cdot\mathcal{D} - \frac{2g_A}{F_\pi}S^\mu \boldsymbol{\tau}\cdot D_\mu\boldsymbol{\pi}\right)N, \tag{63}$$

where $g_A \simeq 1.267$ is the pion-nucleon axial-vector coupling. At this order the nucleon is static; kinetic corrections have relative size $\mathcal{O}(Q/M_{\rm QCD})$ and appear in $\mathcal{L}^{(1)}$.

The formalism to include chiral-symmetry-breaking operators in the $SU(2)\times SU(2)$ χPT Lagrangian has been developed in Refs. [25, 54]. Operators that break the symmetry as components of chiral tensors can be obtained by rotating operators constructed with covariant fields Ψ such as nucleon fields, and nucleon and pion covariant derivatives,

$$\mathcal{O}_{\alpha\beta\cdots\omega}(\boldsymbol{\pi},\Psi) = R_{\alpha\alpha'}(\boldsymbol{\pi})R_{\beta\beta'}(\boldsymbol{\pi})\cdots R_{\omega\omega'}(\boldsymbol{\pi})\mathcal{O}_{\alpha'\beta'\cdots\omega'}(0,\Psi), \tag{64}$$

where the chiral rotation R is given in stereographic coordinates by

$$R_{\alpha\beta}(\boldsymbol{\pi}) = \begin{pmatrix} \delta_{ij} - \frac{2}{D}\frac{\pi_i\pi_j}{F_\pi^2} & \frac{2}{D}\frac{\pi_j}{F_\pi} \\ -\frac{2}{D}\frac{\pi_i}{F_\pi} & \frac{1}{D}\left(1-\frac{\boldsymbol{\pi}^2}{F_\pi^2}\right) \end{pmatrix}. \tag{65}$$

In the EFT these operators generate interactions, now involving $\boldsymbol{\pi}$ directly, that transform as tensors and their tensor products. The strengths of these interactions are proportional to powers of the symmetry-breaking parameters, other dimensional factors being estimated by NDA.

The most important chiral-breaking term is the $\bar{m}$ term in Eq. (38). It generates in the effective Lagrangian all possible interactions with the structures S_4 , $S_4\otimes S_4$, *etc.* with strengths proportional to, respectively, $\bar{m}r(\bar{\theta})$, $(\bar{m}r(\bar{\theta}))^2$, *etc.* In a similar way, one can incorporate in the EFT [54, 56] the $\varepsilon\bar{m}$ term in Eq. (38), which leads to isospin violation as P_3 and its tensor products. The most important terms are, omitting a constant irrelevant for our purposes,

$$\mathcal{L}^{(0,2)}_{\chi,f=0} = -\frac{1}{2D}\left(m_\pi^2 + \frac{\Delta m_\pi^2}{D}\right)\boldsymbol{\pi}^2 + \frac{\delta m_\pi^2}{2D^2}\pi_3^2 \tag{66}$$

and

$$\mathcal{L}^{(1)}_{\chi,f=2} = \Delta m_N\left(1-\frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right)\bar{N}N + \frac{\delta m_N}{2}\bar{N}\left(\tau_3 - \frac{2\pi_3}{F_\pi^2 D}\boldsymbol{\tau}\cdot\boldsymbol{\pi}\right)N. \tag{67}$$

Here the dominant contribution to the pion mass is $m_\pi^2 = \mathcal{O}(r(\bar{\theta})\bar{m}M_{\rm QCD})$, its correction $\Delta m_\pi^2 = \mathcal{O}(m_\pi^4/M_{\rm QCD}^2)$, and the nucleon sigma term $\Delta m_N = \mathcal{O}(m_\pi^2/M_{\rm QCD})$, as the respective interactions have the structures of S_4, $S_4\otimes S_4$, and S_4. In addition, the quark-mass contribution to the nucleon mass splitting is $\delta m_N = \mathcal{O}(r(\bar{\theta})^{-1}\varepsilon\bar{m}) = \mathcal{O}(r(\bar{\theta})^{-2}\varepsilon m_\pi^2/M_{\rm QCD})$ and to the squared pion mass splitting $\delta m_\pi^2 = \mathcal{O}(\delta m_N^2)$, as the respective interactions have the structures of P_3 and $P_3\otimes P_3$. When we are interested in processes with typical momenta $Q\sim m_\pi$, it is convenient to trade $\bar{m}$ by $m_\pi^2/M_{\rm QCD}$ in all chiral-variant terms, which then have strengths proportional to powers of m_π^2 times appropriate powers of $M_{\rm QCD}$. Since $\varepsilon\sim 1/3$ [57], we choose, for simplicity, to count it as $\mathcal{O}(1)$. Since by all evidence $\bar{\theta}$ is small, we also take $r(\bar{\theta})$ as of $\mathcal{O}(1)$.

The power counting of $f \leq 2$ interactions, Eq. (62), is straightforwardly generalized by defining d to count powers of m_π as well [23].

Other isospin-violating hadronic operators and interactions with the photon field A_μ come from $\mathcal{L}_e$, Eq. (34). For applications to EDMs, it is important to construct electromagnetic operators in which hadrons interact with soft photons (with momenta below $M_{\rm QCD}$) in a gauge-invariant way. We can minimally couple charged pions and nucleons to the photon by modifying their covariant derivatives,

$$\begin{aligned} (D_\mu \pi)_i &\rightarrow (D_{\mu,\rm em}\pi)_i = \frac{1}{D}\left(\partial_\mu \delta_{ij} + eA_\mu \epsilon_{3ij}\right)\pi_j, \\ \mathcal{D}_\mu N &\rightarrow \mathcal{D}_{\mu,\rm em} N = \left[\partial_\mu + \frac{i}{F_\pi^2}\boldsymbol{\tau}\cdot\left(\boldsymbol{\pi}\times D_{\mu,\rm em}\boldsymbol{\pi}\right) + \frac{ie}{2}A_\mu\left(1+\tau_3\right)\right] N. \end{aligned} \tag{68}$$

For brevity, in the following we omit the label "em". In addition, we can couple the photon through the field strength $F_{\mu\nu}$, in operators that transform as $I^\mu/6 + T^\mu_{34}/2$ or its tensor products. The index Δ defined in Eq. (62) can be generalized to label electromagnetic operators, by enlarging the definition of d to count also the number of photon fields, which, having dimension one, require compensating powers of $M_{\rm QCD}$ in their coefficients. Finally, integrated-out hard photons give rise to purely hadronic operators that also transform as tensor products of $I^\mu/6 + T^\mu_{34}/2$ [54], and are proportional to powers of $(e/4\pi)^2 = \alpha_{\rm em}/4\pi$. The most important of these operators is the electromagnetic contribution $\breve{\delta} m_\pi^2$ to the squared pion mass splitting,

$$\mathcal{L}^{(1)}_{{\rm em},f=0} = -\frac{\breve{\delta} m_\pi^2}{2D^2}\left(\boldsymbol{\pi}^2 - \pi_3^2\right), \tag{69}$$

with $\breve{\delta} m_\pi^2 = \mathcal{O}(\alpha_{\rm em} M_{\rm QCD}^2/4\pi)$. Since numerically $\alpha_{\rm em}/4\pi \simeq (m_\pi/M_{\rm QCD})^3$, we assign $d = 3$ for each power of $\alpha_{\rm em}/4\pi$ [54]. This counting reflects both the smallness and the presumed electromagnetic origin of the pion mass splitting.

The remaining dimension-four term, the $m_* \sin\bar\theta$ term in Eq. (38), which is $\not{P}\not{T}$, can be treated in the same way. This was done in some detail in Ref. [32], where the implications of its P_4 structure were discussed. Perhaps the most important point is that the hadronic $\not{P}\not{T}$ interactions are intrinsically linked to isospin-breaking operators that transform like P_3, so that the $\not{P}\not{T}$ LECs can be inferred from isospin-breaking couplings, when the latter are known.

In the rest of this paper we discuss the construction of the $\not{P}\not{T}$ interactions stemming from $\mathcal{L}_6$, Eq. (54), and their power counting. The $\not{P}\not{T}$ chiral Lagrangian can be constructed by writing down all terms that transform in the same way under Lorentz, P, T, and chiral symmetry as the terms in Eq. (54). Because such interactions are very small compared to CP-even interactions, we restrict ourselves to low-energy operators involving at most one power of the dimensionless coefficients $\delta_{0,3}$, $\tilde{\delta}_{0,3}$, w, ξ, and $\sigma_{1,8}$, even though it is straightforward to consider mixed effects. As discussed above, these coefficients are model dependent, and we construct the low-energy $\not{P}\not{T}$ Lagrangian for each source separately. The form of the low-energy interactions is determined by the way a source breaks chiral symmetry.

The gCEDM and the two four-quark operators in Eq. (54) with coefficients ${\rm Im}\Sigma_{1,8}$ do not break chiral symmetry and are therefore $SO(4)$ scalars, which implies that they induce the same chiral Lagrangian and cannot be separated on the basis of low-energy experiments alone. From now on we refer to these as chiral-invariant sources (χISs) and use the symbol w ($\bar{I}$) to denote collectively the dimensionless constants w and $\sigma_{1,8}$ (the invariants I_W, $I^{(1)}_{qq}$, and $I^{(8)}_{qq}$) in Eq. (48)

(Eq. (49)):

$$\{w, \sigma_1, \sigma_8\} \;\; \to \;\; w, \tag{70}$$

$$\left\{I_W, I_{qq}^{(1)}, I_{qq}^{(8)}\right\} \;\; \to \;\; \bar{I}. \tag{71}$$

Pion interactions stemming from χISs involve the pion covariant derivative, unless $\bar{I}$ is in a tensor product with a chiral-variant tensor (*i.e.*, a product of Ss and Ps).

In contrast to the χISs, the other sources (qEDM, qCEDM, FQLR) break chiral symmetry explicitly and can bring in pion non-derivative interactions by themselves, as is the case for $\bar{\theta}$. The specific dependence on $\boldsymbol{\pi}$ depends on the source. The qCEDM consists of two independent components, the isoscalar and isovector qCEDMs. They transform, respectively, as the fourth and third components of the $SO(4)$ vectors $\tilde{V}$ and $\tilde{W}$ defined in Eq. (51). Since the QCD $\bar{\theta}$ term transforms as the fourth component of an $SO(4)$ vector as well, the $\bar{\theta}$ term and isoscalar qCEDM generate identical chiral operators (but with different strenghts).

By themselves, the FQLR operators have the most complicated chiral structure, transforming as the 34-component of the symmetric tensors $X^{(1)}$ and $X^{(8)}$ in Eqs. (52) and (53). As for χISs, both operators induce the same chiral Lagrangian, but their entanglement is not a real issue since they depend on the same dimensionless parameter ξ. From now on we use the symbol X to denote $X^{(1)}$ and $X^{(8)}$:

$$\left\{X^{(1)}, X^{(8)}\right\} \;\; \to \;\; X. \tag{72}$$

The structure of the qEDM, transforming as the fourth and third components of the $SO(4)$ vectors V and W in Eq. (50), resembles that of the qCEDM, but the similarity is deceiving due to the photon. When the photon is integrated out to produce purely hadronic operators from qEDM, one needs to take the tensor product with the $I^\mu/6 + T_{34}^\mu/2$ from Eq. (34). Conversely, to produce operators with a soft-photon the qCEDM requires (as do FQLR and χISs) a tensor product with $I^\mu/6 + T_{34}^\mu/2$. In both cases, the extra T_{34}^μ produces in general interactions of more complicated structure.

Note that for each chiral-variant $\not{P}\not{T}$ source, there are associated quark-gluon operators that do not violate P and T. For example, a qEDM is associated with a quark magnetic dipole moment, since they make the same $SO(4)$ vector, Eq. (50). Analogous statements hold for qCEDM in Eq. (51) and for FQLR in Eqs. (52) and (53). The situation is similar to the $\bar{\theta}$ term, which is related to the quark mass splitting [32]. However, for dimension-six sources these relations are less useful, since the corresponding CP-even interactions are buried among much larger, dimension-four interactions.

The interactions stemming from the dimension-six $\not{P}\not{T}$ sources can be organized according to a chiral index analogous to Eq. (62), with the only difference that the coefficients of low-energy interactions must contain two inverse powers of the high-energy scale $M_{\not{T}}$, which replace two inverse powers of M_{QCD}. The powers of M_{QCD} in a coefficient are therefore counted by

$$\Delta_6 = d + f/2 - 4, \tag{73}$$

where d is the number of derivatives, and powers of the quark mass and e as described above. We will find that for the qEDM, qCEDM and χISs $\Delta_6 \geq -2$, while for the FQLR operator $\Delta_6 \geq -4$. The suppression by at least one power of $\alpha_{\text{em}}/4\pi$ renders the purely hadronic operators from hard photons essentially irrelevant.

The question now is to what order, for each separate source, do we need to construct the $\not{P}\not{T}$ chiral Lagrangian. This obviously depends on what observable one wants to calculate. The most likely candidates are $\not{P}\not{T}$ electromagnetic moments and form factors of light nuclei, which require the use of the most important $\not{P}\not{T}$ pion, pion-nucleon, photon-nucleon, and inter-nucleon interactions. In order to compare our EFT approach to more traditional approaches, where $\not{P}\not{T}$ is implemented through three non-derivative $\not{P}\not{T}$ pion-nucleon interactions, we construct the pion-nucleon Lagrangian for each source up to the order where all three pion-nucleon interactions appear. As found in Refs. [35, 36], for most sources the $\not{P}\not{T}$ nucleon-nucleon(-photon) interactions are subleading with respect to $\not{P}\not{T}$ one-pion exchange, so in the nucleon-nucleon sector we only construct LO operators. We construct the electromagnetic sector until, for each source, we find the momentum dependence of the nucleon EDFF, as the linear term in the square of the momentum transferred is a contribution the Schiff moment, which is important for the evaluation of atomic EDMs [28]. Since operators with two explicit photons give small contributions even to atomic EDMs [58], we do not construct explicitly here operators with more than a single soft photon.

We start in Sec. 5 with the construction of operators in the purely mesonic sector (that is, with $f = 0$). We then tackle the pion- and photon-nucleon sectors ($f = 2$) in Secs. 6 and 8, respectively. We extend our analysis to include the nuclear sector in Sec. 10.

5 Pion sector

In contrast to CP-even dynamics, $\not{P}\not{T}$ can generate interactions with an odd number of pions, starting with pion tadpoles that represent the disappearance of a neutral pion into the vacuum. Tadpoles cause the vacuum to become unstable because it can create neutral pions to lower its energy. In the $\bar{\theta}$ case, imposing the condition of vacuum alignment at the quark level in first order in the quark masses [44] ensures that the $\not{P}\not{T}$ breaking is through the $SO(4)$ vector P in Eq. (39) and that tadpoles are absent up to that order. For the dimension-six sources, the more complicated chiral structure also leads to tadpoles. Since the tadpoles are small, they can, in principle, be treated in perturbation theory, but a more convenient way of handling them is to rotate them away by performing a field redefinition [32]. After we consider the two leading orders for each source, we discuss the effects of tadpole extermination, which has important consequences for other sectors of the theory.

5.1 qCEDM

As it was found in Ref. [32], without any nucleon fields it is not possible to construct an operator that transforms as the fourth component of an $SO(4)$ vector, so the isoscalar qCEDM has no effect without another source of chiral breaking. The isovector qCEDM, however, breaks isospin symmetry as $\tilde{d}_3\tilde{W}_3$ and generates a pion tadpole with chiral index $\Delta_6 = -2$,

$$\mathcal{L}^{(-2)}_{\tilde{q},f=0} = \bar{\Delta}^{(-2)}_{\tilde{q}}\frac{F_\pi\pi_3}{2D}, \tag{74}$$

where, by NDA, the LEC scales as

$$\bar{\Delta}^{(-2)}_{\tilde{q}} = \mathcal{O}\left(\tilde{\delta}_3\frac{m_\pi^2 M_{\text{QCD}}^2}{M_{\not{T}}^2}\right). \tag{75}$$

The next operators appear two orders higher in the chiral expansion. First, there is a $\tilde{d}_3\tilde{W}_3$-type operator built from two pion covariant derivatives. Second, there are contributions from the combined effect of the quark mass and the qCEDM, producing pionic operators with the same chiral properties as the tensor products $\tilde{d}_3\tilde{W}_3 \otimes \bar{m}S_4$ and $\tilde{d}_0\tilde{V}_4 \otimes \bar{m}\varepsilon P_3$. Together, these are

$$\mathcal{L}^{(0)}_{\tilde{q},f=0} = \vartheta_1 \frac{\pi_3}{F_\pi D}\left(D_\mu \boldsymbol{\pi} \cdot D^\mu \boldsymbol{\pi}\right) + \bar{\Delta}^{(0)}_{\tilde{q}} \frac{F_\pi \pi_3}{2D}\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right), \tag{76}$$

where

$$\vartheta_1 = \mathcal{O}\left(\tilde{\delta}_3 \frac{m_\pi^2}{M_{\not{T}}^2}\right), \qquad \bar{\Delta}^{(0)}_{\tilde{q}} = \mathcal{O}\left((\varepsilon\tilde{\delta}_0 + \tilde{\delta}_3)\frac{m_\pi^4}{M_{\not{T}}^2}\right). \tag{77}$$

The $+$ in the scaling of $\bar{\Delta}^{(0)}_{\tilde{q}}$ should not be taken literally, but as an indication that the LEC gets contributions from two sources.

5.2 FQLR

In the pionic sector, X_{34} also leads to a tadpole operator, but of different form than qCEDM's:

$$\mathcal{L}^{(-4)}_{\mathrm{LR},f=0} = \bar{\Delta}^{(-4)}_{\mathrm{LR}} \frac{F_\pi \pi_3}{2D}\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right), \tag{78}$$

with coefficient

$$\bar{\Delta}^{(-4)}_{\mathrm{LR}} = \mathcal{O}\left(\xi \frac{M_{\mathrm{QCD}}^4}{M_{\not{T}}^2}\right). \tag{79}$$

Equation (78) generates different three-pion vertices than Eq. (74). As we will see, this difference is important, because the elimination of the pion tadpole does not completely cancel the operator in Eq. (78); instead, it leaves some three-pion coupling behind.

Two orders down we find terms with two derivatives and terms transforming as $\mathrm{Im}\,\Xi\, X_{34} \otimes \bar{m}S_4$,

$$\begin{aligned}\mathcal{L}^{(-2)}_{\mathrm{LR},f=0} \;=\;& \vartheta_2 \frac{\pi_3}{F_\pi D^2}\left(D_\mu \boldsymbol{\pi} \cdot D^\mu \boldsymbol{\pi}\right)\left(1 - \frac{\boldsymbol{\pi}^2}{F_\pi^2}\right) + \vartheta_3 \frac{\boldsymbol{\pi}\cdot D_\mu \boldsymbol{\pi}}{F_\pi D}\left(D^\mu \pi_3 - \frac{2}{F_\pi^2 D}\pi_3 \boldsymbol{\pi}\cdot D^\mu \boldsymbol{\pi}\right) \\ &+ \frac{F_\pi \pi_3}{2D}\left[\bar{\Delta}^{(-2)}_{\mathrm{LR1}} + \bar{\Delta}^{(-2)}_{\mathrm{LR2}}\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right)^2\right],\end{aligned} \tag{80}$$

with the scalings

$$\vartheta_{2,3} = \mathcal{O}\left(\xi \frac{M_{\mathrm{QCD}}^2}{M_{\not{T}}^2}\right), \qquad \bar{\Delta}^{(-2)}_{\mathrm{LR1,2}} = \mathcal{O}\left(\xi \frac{m_\pi^2 M_{\mathrm{QCD}}^2}{M_{\not{T}}^2}\right). \tag{81}$$

5.3 qEDM

The leading pion operators induced by the qEDM transform as $d_3W_3 \otimes eI^\mu/6$ or $d_0V_4 \otimes eT^\mu_{34}/2$, and are simply given by

$$\mathcal{L}^{(1)}_{q,f=0} = \bar{\Delta}^{(1)}_{q} \frac{F_\pi \pi_3}{2D}, \tag{82}$$

where the LEC scales as

$$\bar{\Delta}_q^{(1)} = \mathcal{O}\left(\frac{\alpha_{\rm em}}{4\pi}(\delta_0+\delta_3)\frac{m_\pi^2 M_{\rm QCD}^2}{M_{\not{T}}^2}\right). \tag{83}$$

Again the + in the scaling of the LEC should not be taken literally. Because of the $\alpha_{\rm em}/4\pi$ suppression, we do not bother to go to higher order.

5.4 χISs

In the mesonic sector it is not possible to write down $\not{P}\not{T}$ chiral-invariant operators. Operators in this sector can be constructed by combining the effects of χISs with chiral-symmetry breaking from the quark mass difference, which renders them identical in form to those of the qCEDM.

Indeed, at lowest order $d_W \bar{I} \otimes \varepsilon \bar{m} P_3$ gives

$$\mathcal{L}_{w,f=0}^{(-2)} = \bar{\Delta}_w^{(-2)} \frac{F_\pi \pi_3}{2D}, \tag{84}$$

with

$$\bar{\Delta}_w^{(-2)} = \mathcal{O}\left(w\varepsilon \frac{m_\pi^2 M_{\rm QCD}^2}{M_{\not{T}}^2}\right). \tag{85}$$

At $\Delta_6 = 0$, we get operators with two covariant derivatives and operators transforming as $d_W \bar{I} \otimes \bar{m} S_4 \otimes \bar{m}\varepsilon P_3$,

$$\mathcal{L}_{w,f=0}^{(0)} = \vartheta_1 \frac{\pi_3}{F_\pi D}\left(D_\mu \boldsymbol{\pi}\cdot D^\mu \boldsymbol{\pi}\right) + \bar{\Delta}_w^{(0)} \frac{\pi_3 F_\pi}{2D}\left(1-\frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right), \tag{86}$$

where the LECs scale as

$$\vartheta_1 = \mathcal{O}\left(w\varepsilon \frac{m_\pi^2}{M_{\not{T}}^2}\right), \qquad \bar{\Delta}_w^{(0)} = \mathcal{O}\left(w\varepsilon \frac{m_\pi^4}{M_{\not{T}}^2}\right). \tag{87}$$

5.5 Tadpole extermination

We have just seen that the transformation properties of the $\not{P}\not{T}$ dimension-six sources cause pion tadpoles to appear in the $\not{P}\not{T}$ mesonic Lagrangian. Since the coupling constant of the neutral pion to the vacuum is small compared to the pion mass, these tadpoles can be dealt with in perturbation theory, meaning that, for any given $\not{P}\not{T}$ observable at a given accuracy, only a finite number of neutral pions disappearing into the vacuum must be considered [32]. For applications, such as the calculation of hadronic [43] and nuclear [34, 36] EDMs, it is however more convenient to eliminate the pion tadpoles from the mesonic Lagrangian, which can be achieved with field redefinitions of the form discussed in Ref. [32].

We define a new pion field $\boldsymbol{\zeta}' = \boldsymbol{\pi}'/F_\pi$ through

$$\zeta_i = \frac{1}{d'}\left\{\zeta_i' - \delta_{i3}[2C\zeta_3' + S(1-\boldsymbol{\zeta}'^2)]\right\}, \tag{88}$$

where

$$d' = 1 - C(1-\boldsymbol{\zeta}'^2) + 2S\zeta_3', \tag{89}$$

and

$$C = \frac{1}{2}(1 - \cos\varphi), \qquad S = \frac{1}{2}\sin\varphi, \tag{90}$$

in terms of an angle φ. This transformation looks complicated, but has the nice property that the pion covariant derivative transforms simply as

$$D_\mu \zeta_i = O'_{ij} D'_\mu \zeta'_j, \tag{91}$$

with an orthogonal matrix

$$O'_{ij} = \delta_{ij} - \frac{2}{d'} \left\{ C \left[(\boldsymbol{\zeta}'^2 - \zeta_3'^2)\delta_{ij} - \epsilon_{3ik}\epsilon_{3jl}\zeta'_k\zeta'_l \right] + (C\zeta'_3 + S)\left(\zeta'_i\delta_{3j} - \zeta'_j\delta_{3i}\right) \right\}. \tag{92}$$

This simple transformation ensures that chiral-invariant operators, built from pion covariant derivatives, are invariant under the field redefinition in Eq. (88). This is not the case for operators that break chiral symmetry and involve the pion field directly.

Before performing any rotation we summarize here the non-derivative, chiral-symmetry-breaking Lagrangian in the purely mesonic sector, including quark mass and $\not{P}\not{T}$ operators in the first two orders (except for the very small qEDM operators, for which we keep only the least unimportant term). From Eqs. (66), (74), (76), (78), (80), (82), (84), and (86),

$$\begin{aligned}
\mathcal{L}_{\text{tadpole}} &= -\frac{m_\pi^2}{2D}\left(1 + \frac{\Delta m_\pi^2}{m_\pi^2 D}\right)\boldsymbol{\pi}^2 + \frac{\delta m_\pi^2}{2D^2}\pi_3^2 + \left(\bar{\Delta}_{\text{LR1}}^{(-2)} + \bar{\Delta}_{\tilde{q}}^{(-2)} + \bar{\Delta}_q^{(1)} + \bar{\Delta}_w^{(-2)}\right)\frac{F_\pi\pi_3}{2D} \\
&\quad + \left(\bar{\Delta}_{\text{LR}}^{(-4)} + \bar{\Delta}_{\tilde{q}}^{(0)} + \bar{\Delta}_w^{(0)}\right)\frac{F_\pi\pi_3}{2D}\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right) + \bar{\Delta}_{\text{LR2}}^{(-2)}\frac{F_\pi\pi_3}{2D}\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right)^2.
\end{aligned} \tag{93}$$

We omit chiral-breaking operators generated by the electromagnetic interaction, which are not affected by the field redefinitions [32].

Our first goal is to remove for each source the dominant tadpole, that is, the terms with index $\Delta_6 = -4$ for FQLR, $\Delta_6 = -2$ for qCEDM and χISs, and $\Delta_6 = 1$ for qEDM. By performing the field redefinitions in Eq. (88) with the small angle

$$\begin{aligned}
\varphi \simeq \tan\varphi &= -\frac{1}{m_\pi^2}\left(\bar{\Delta}_{\text{LR}}^{(-4)} + \bar{\Delta}_{\tilde{q}}^{(-2)} + \bar{\Delta}_w^{(-2)} + \bar{\Delta}_q^{(1)}\right) \\
&= \mathcal{O}\left(\left(\xi\frac{M_{\text{QCD}}^2}{m_\pi^2} + \tilde{\delta}_3 + w\varepsilon + \frac{\alpha_{\text{em}}}{4\pi}(\delta_0 + \delta_3)\right)\frac{M_{\text{QCD}}^2}{M_{\not{T}}^2}\right),
\end{aligned} \tag{94}$$

the dominant tadpoles are removed. From here on we keep only terms linear in φ. The effect of this field redefinition on the mesonic Lagrangian, apart from canceling the dominant tadpoles, is to modify the coefficients of the tensors:

$$\begin{aligned}
\mathcal{L}_{\text{tadpole}} &= -\frac{m_\pi^2}{2D}\left(1 + \frac{\Delta m_\pi^2}{m_\pi^2 D}\right)\boldsymbol{\pi}^2 + \frac{\delta m_\pi^2}{2D^2}\pi_3^2 - \bar{\Delta}_{\text{LR}}^{(-4)}\frac{\pi_3\boldsymbol{\pi}^2}{F_\pi D^2} + \bar{\Delta}_{\text{LR1}}^{(-2)}\frac{F_\pi\pi_3}{2D} \\
&\quad + \left(\bar{\Delta}_{\text{LR}}^{(-2)\prime} + \bar{\Delta}_{\tilde{q}}^{(0)\prime} + \bar{\Delta}_w^{(0)\prime}\right)\frac{F_\pi\pi_3}{2D}\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right) + \bar{\Delta}_{\text{LR2}}^{(-2)}\frac{F_\pi\pi_3}{2D}\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right)^2,
\end{aligned} \tag{95}$$

in terms of the shifted LECs,

$$
\begin{aligned}
\bar{\Delta}^{(-2)\prime}_{\mathrm{LR}} &= -\frac{\Delta m_\pi^2 - \delta m_\pi^2}{m_\pi^2}\,\bar{\Delta}^{(-4)}_{\mathrm{LR}} = \mathcal{O}\left(\xi(1+\varepsilon^2)\frac{m_\pi^2 M_{\mathrm{QCD}}^2}{M_{\not{T}}^2}\right), &(96)\\
\bar{\Delta}^{(0)\prime}_{\tilde{q}} &= \bar{\Delta}^{(0)}_{\tilde{q}} - \frac{\Delta m_\pi^2 - \delta m_\pi^2}{m_\pi^2}\,\bar{\Delta}^{(-2)}_{\tilde{q}} = \mathcal{O}\left(\left(\varepsilon\tilde{\delta}_0 + (1+\varepsilon^2)\tilde{\delta}_3\right)\frac{m_\pi^4}{M_{\not{T}}^2}\right), &(97)\\
\bar{\Delta}^{(0)\prime}_{w} &= \bar{\Delta}^{(0)}_{w} - \frac{\Delta m_\pi^2 - \delta m_\pi^2}{m_\pi^2}\,\bar{\Delta}^{(-2)}_{w} = \mathcal{O}\left(w\varepsilon(1+\varepsilon^2)\frac{m_\pi^4}{M_{\not{T}}^2}\right). &(98)
\end{aligned}
$$

(Here and in the following we neglect terms of higher order than we need in the rest of the paper.) We can thus rotate away the leading tadpoles without introducing new interactions in the meson Lagrangian. The net effect of the rotation is only to change the dependence of the coefficients on the parameters $\tilde{\delta}_0$, $\tilde{\delta}_3$, and ε.

The remaining tadpoles in Eq. (95) can be eliminated by a second rotation, now with an even smaller angle

$$
\varphi' = -\frac{1}{m_\pi^2}\left(\bar{\Delta}^{(-2)}_{\mathrm{LR1}} + \bar{\Delta}^{(-2)}_{\mathrm{LR2}} + \bar{\Delta}^{(-2)\prime}_{\mathrm{LR}} + \bar{\Delta}^{(0)\prime}_{\tilde{q}} + \bar{\Delta}^{(0)\prime}_{w}\right), \tag{99}
$$

which leaves us with

$$
\begin{aligned}
\mathcal{L}_{\mathrm{tadpole}} &= -\frac{m_\pi^2}{2D}\left(1 + \frac{\Delta m_\pi^2}{m_\pi^2 D}\right)\boldsymbol{\pi}^2 + \frac{\delta m_\pi^2}{2D^2}\pi_3^2\\
&\quad -\frac{\pi_3\boldsymbol{\pi}^2}{F_\pi D^2}\left(\bar{\Delta}^{(-4)}_{\mathrm{LR}} + \bar{\Delta}^{(-2)\prime}_{\mathrm{LR}} + \frac{2\bar{\Delta}^{(-2)}_{\mathrm{LR2}}}{D} + \bar{\Delta}^{(0)\prime}_{\tilde{q}} + \bar{\Delta}^{(0)\prime}_{w}\right). &(100)
\end{aligned}
$$

Although the tadpoles are removed, residual $\not{P}\not{T}$ interactions involving an odd number of pions are left behind. In case of the qCEDM and χISs these terms carry a high chiral index and such multi-pion vertices contribute to observables at high order. In case of the FQLR, due to its complicated $SO(4)$ properties, a three-pion interaction remains with a relatively low index. The consequences of this interaction are discussed in Secs. 7.4, 9, and 10.5.

So far we did not discuss the operators with LECs ϑ_i. These operators are not affected by the above rotations up to the order we are working. They induce changes in the pion kinetic term, which nevertheless can be removed by another pion-field redefinition.

6 Pion-nucleon sector

For most applications, we can ignore $\not{P}\not{T}$ effects in the pion sector, once the tadpoles have been removed. Tadpole extermination does leave traces, however, in operators involving nucleons. In this section we construct the $\not{P}\not{T}$ pion-nucleon operators, incorporating the new terms arising from the field redefinitions needed to remove the tadpoles. $\not{P}\not{T}$ pion-nucleon interactions are very important for nucleon and nuclear EDMs.

6.1 Tadpoles strike back

It is extremely convenient to supplement the pion field redefinition (88) with a nucleon field redefinition [32]

$$N = U'N', \tag{101}$$

in terms of the unitary matrix

$$U' = \frac{1}{\sqrt{d'}}\left[\sqrt{1-C}+\sqrt{C}(\zeta'_3 + i\epsilon_{3jk}\zeta'_j\tau_k)\right], \tag{102}$$

where d', C, and S are given by Eqs. (89) and (90). This redefinition ensures simple transformation laws for covariant objects such as nucleon covariant derivatives and nucleon bilinears,

$$\mathcal{D}_\mu N = U'\mathcal{D}'_\mu N', \qquad \bar{N}N = \bar{N}'N', \qquad \bar{N}\tau_i N = O'_{ij}\bar{N}'\tau_j N', \tag{103}$$

where O' is the orthogonal matrix in Eq. (92). The properties (103) extend to the nucleon sector the invariance under field redefinition of chiral-invariant operators, built from nucleon fields and covariant derivatives.

The two rotations that allow elimination of the pion tadpoles in leading orders affect the chiral-breaking terms involving nucleons. Most importantly, the nucleon sigma term and mass splitting in Eq. (67) become, after the rotation of Eq. (94),

$$\mathcal{L}_{\not\chi,f=2} = \Delta m_N\left(1-\frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}+\varphi\frac{2\pi_3}{F_\pi D}\right)\bar{N}N+\frac{\delta m_N}{2}\bar{N}\left(\tau_3-\frac{2\pi_3}{F_\pi^2 D}\boldsymbol{\tau}\cdot\boldsymbol{\pi}+\varphi\frac{2}{F_\pi D}\boldsymbol{\tau}\cdot\boldsymbol{\pi}\right)N. \tag{104}$$

The terms linear in φ contribute to the $\not{P}\not{T}$ pion-nucleon interactions, in addition to those generated directly from the chiral structure of the $\not{P}\not{T}$ dimension-six sources. Higher-order pion-nucleon operators come from rotation with angle φ' (99) and/or higher-order chiral-breaking terms in the CP-even Lagrangian.

We refrain from giving the boring details here. Instead, we turn to the construction of the $\not{P}\not{T}$ pion-nucleon interactions from the chiral structure of the various sources. We indicate below the changes that arise from tadpole extermination.

6.2 qCEDM

Since the qCEDM breaks chiral symmetry, it gives rise to non-derivative pion-nucleon interactions. Already at chiral order $\Delta_6 = -1$, the structures $\tilde{d}_0\tilde{V}_4$ and $\tilde{d}_3\tilde{W}_3$ give

$$\mathcal{L}^{(-1)}_{\tilde{q},f=2} = -\frac{\bar{g}_0}{F_\pi D}\bar{N}\boldsymbol{\tau}\cdot\boldsymbol{\pi}N-\frac{\bar{g}_1}{F_\pi D}\pi_3\bar{N}N, \tag{105}$$

with

$$\bar{g}_0 = \mathcal{O}\left(\tilde{\delta}_0\frac{m_\pi^2 M_{\mathrm{QCD}}}{M_{\not{T}}^2}\right), \qquad \bar{g}_1 = \mathcal{O}\left(\tilde{\delta}_3\frac{m_\pi^2 M_{\mathrm{QCD}}}{M_{\not{T}}^2}\right). \tag{106}$$

Comparing Eqs. (105) and (104) shows that tadpole removal leads to shifts

$$\bar{g}_0 \;\to\; \bar{g}_0+\delta m_N\frac{\bar{\Delta}^{(-2)}_{\tilde{q}}}{m_\pi^2} = \mathcal{O}\left(\left(\tilde{\delta}_0+\varepsilon\tilde{\delta}_3\right)\frac{m_\pi^2 M_{\mathrm{QCD}}}{M_{\not{T}}^2}\right), \tag{107}$$

$$\bar{g}_1 \;\to\; \bar{g}_1+2\Delta m_N\frac{\bar{\Delta}^{(-2)}_{\tilde{q}}}{m_\pi^2} = \mathcal{O}\left(\tilde{\delta}_3\frac{m_\pi^2 M_{\mathrm{QCD}}}{M_{\not{T}}^2}\right). \tag{108}$$

While $\bar{g}_1$ gets a change of $\mathcal{O}(1)$ that nevertheless does not affect its scaling, $\bar{g}_0$ now depends on the isovector qCEDM as well. In the following we absorb the terms $\propto \bar{\Delta}_{\tilde{q}}^{(-2)}$ in $\bar{g}_i$.

Equation (105) shows the first important difference between $\not{P}\not{T}$ from dimension-six operators and the QCD $\bar{\theta}$ term, namely, the presence of the $\not{P}\not{T}$ isospin-breaking interaction with LEC $\bar{g}_1$ at leading order in the $f = 2$ Lagrangian. This fact is particularly relevant for the $\not{P}\not{T}$ moments of the deuteron [59, 34, 36]. The $\bar{\theta}$ term also generates this interaction but it is suppressed by $(m_\pi/M_{\text{QCD}})^2$ compared to leading-order interactions [32].

Increasing Δ_6 by one, we find operators with the same chiral structures but one covariant derivative,

$$\begin{aligned}\mathcal{L}_{\tilde{q},f=2}^{(0)} &= \frac{2\bar{\beta}_1}{F_\pi^2 D}(\boldsymbol{\pi}\cdot\mathcal{D}_\mu\boldsymbol{\pi})\bar{N}S^\mu N + \frac{2\bar{\beta}_2}{F_\pi^2 D}(\pi_3\mathcal{D}_\mu\boldsymbol{\pi})\cdot\bar{N}S^\mu\boldsymbol{\tau}N \\ &\quad +\frac{\bar{\beta}_3}{F_\pi}\left(\delta_{i3}-\frac{2\pi_i\pi_3}{F_\pi^2 D}\right)\bar{N}(\boldsymbol{\tau}\times v\cdot D\boldsymbol{\pi})_i N,\end{aligned}\tag{109}$$

where the LECs are

$$\bar{\beta}_1 = \mathcal{O}\left(\tilde{\delta}_0\frac{m_\pi^2}{M_{\not{T}}^2}\right),\qquad \bar{\beta}_{2,3} = \mathcal{O}\left(\tilde{\delta}_3\frac{m_\pi^2}{M_{\not{T}}^2}\right).\tag{110}$$

Increasing Δ_6 by another unit, the chiral structures proliferate significantly, and so does the number of different interactions. With still the same chiral structure as before, but two covariant derivatives, we find

$$\begin{aligned}\mathcal{L}_{\tilde{q},f=2}^{(1)} &= \left\{\frac{\bar{\zeta}_1}{F_\pi}(\mathcal{D}_{\mu,\perp}D_\perp^\mu\boldsymbol{\pi})\cdot\bar{N}\boldsymbol{\tau}N + \frac{\bar{\zeta}_2}{F_\pi^2}(D_\mu\boldsymbol{\pi}\times v\cdot D\boldsymbol{\pi})\cdot\bar{N}S^\mu\boldsymbol{\tau}N\right. \\ &\quad \left.+\frac{\bar{\zeta}_3}{F_\pi}(v\cdot\mathcal{D}v\cdot D\boldsymbol{\pi})\cdot\bar{N}\boldsymbol{\tau}N + \frac{\bar{\zeta}_4}{2F_\pi}(D_\nu\boldsymbol{\pi})\cdot\bar{N}[S^\mu,S^\nu]\boldsymbol{\tau}\mathcal{D}_{\mu,-}N\right\}\frac{1}{D}\left(1-\frac{\boldsymbol{\pi}^2}{F_\pi^2}\right) \\ &\quad +\left\{\frac{\bar{\zeta}_5}{4}\bar{N}\boldsymbol{\tau}(v\cdot\mathcal{D}_-)^2N + \frac{\bar{\zeta}_6}{4}\bar{N}\boldsymbol{\tau}\mathcal{D}_{\perp,-}^2N - \frac{\bar{\zeta}_7}{F_\pi}\bar{N}S^\mu[\boldsymbol{\tau}\times(v\cdot\mathcal{D}D_\mu\boldsymbol{\pi})]N\right. \\ &\quad -\frac{i\bar{\zeta}_8}{F_\pi}(v\cdot D\boldsymbol{\pi})\bar{N}S\cdot\mathcal{D}_-N - \frac{2i\bar{\zeta}_9}{F_\pi^2}(D_\mu\boldsymbol{\pi}\times D_\nu\boldsymbol{\pi})\bar{N}[S^\mu,S^\nu]N \\ &\quad -\frac{\bar{\zeta}_{10}}{F_\pi^2}(D_{\mu,\perp}\boldsymbol{\pi})(D_\perp^\mu\boldsymbol{\pi})\cdot\bar{N}\boldsymbol{\tau}N - \frac{\bar{\zeta}_{11}}{F_\pi^2}(v\cdot D\boldsymbol{\pi})(v\cdot D\boldsymbol{\pi})\cdot\bar{N}\boldsymbol{\tau}N \\ &\quad \left.-\frac{\bar{\zeta}_{12}}{F_\pi^2}(D_{\mu,\perp}\boldsymbol{\pi})^2\bar{N}\boldsymbol{\tau}N - \frac{\bar{\zeta}_{13}}{F_\pi^2}(v\cdot D\boldsymbol{\pi})^2\bar{N}\boldsymbol{\tau}N\right\}\cdot\frac{\boldsymbol{\pi}}{F_\pi D} \\ &\quad +\left\{\frac{\bar{\xi}_1}{F_\pi}(\mathcal{D}_{\mu,\perp}D_\perp^\mu\pi_i)\bar{N}N + \frac{\bar{\xi}_2}{F_\pi^2}(D_\mu\boldsymbol{\pi}\times v\cdot D\boldsymbol{\pi})_i\bar{N}S^\mu N + \frac{\bar{\xi}_3}{F_\pi}(v\cdot\mathcal{D}v\cdot D\pi_i)\bar{N}N\right. \\ &\quad \left.+\frac{\bar{\xi}_4}{F_\pi}(D_\nu\pi_i)\bar{N}[S^\mu,S^\nu]\mathcal{D}_{\mu,-}N + \frac{i\bar{\xi}_5}{F_\pi}\bar{N}((D_\mu\boldsymbol{\pi})\times\boldsymbol{\tau})_i\mathcal{D}_{\perp,-}^\mu N\right\}\left(\delta_{i3}-\frac{2\pi_3\pi_i}{F_\pi^2 D}\right) \\ &\quad +\left\{\frac{\bar{\xi}_6}{4}\bar{N}(v\cdot\mathcal{D}_-)^2N + \frac{\bar{\xi}_7}{4}\bar{N}\mathcal{D}_{\perp,-}^2N + \frac{i\bar{\xi}_8}{2F_\pi}(v\cdot D\boldsymbol{\pi})\cdot\bar{N}\boldsymbol{\tau}S\cdot\mathcal{D}_-N\right. \\ &\quad \left.+\frac{i\bar{\xi}_9}{2F_\pi^2}(D_\mu\boldsymbol{\pi}\times D_\nu\boldsymbol{\pi})\cdot\bar{N}[S^\mu,S^\nu]\boldsymbol{\tau}N + \frac{\bar{\xi}_{10}}{F_\pi^2}(D_{\mu,\perp}\boldsymbol{\pi})^2\bar{N}N + \frac{\bar{\xi}_{11}}{F_\pi^2}(v\cdot D\boldsymbol{\pi})^2\bar{N}N\right\}\frac{2\pi_3}{F_\pi D}.\end{aligned}\tag{111}$$

The scaling of the LECs $\bar{\zeta}_i$ and $\bar{\xi}_i$ is given by

$$\bar{\zeta}_i = \mathcal{O}\left(\tilde{\delta}_0 \frac{m_\pi^2}{M_{\not T}^2 M_{\rm QCD}}\right), \qquad \bar{\xi}_i = \mathcal{O}\left(\tilde{\delta}_3 \frac{m_\pi^2}{M_{\not T}^2 M_{\rm QCD}}\right). \tag{112}$$

Reparametrization invariance relates some of these operators to operators of lower order,

$$\begin{aligned} &\bar{\zeta}_4 = \bar{\zeta}_6 = \frac{\bar{g}_0}{2m_N^2}, \qquad \bar{\zeta}_8 = \frac{g_A \bar{g}_0}{m_N^2} - \frac{\bar{\beta}_1}{m_N}, \\ &\bar{\xi}_4 = \bar{\xi}_7 = \frac{\bar{g}_1}{4m_N^2}, \qquad \bar{\xi}_5 = -\frac{\bar{\beta}_3}{2m_N}, \qquad \bar{\xi}_8 = -\frac{g_A \bar{g}_1}{m_N^2} - \frac{\bar{\beta}_2}{m_N}. \end{aligned} \tag{113}$$

In these subleading pion-nucleon Lagrangians, the effects of the field redefinition on operators that transform as $\tilde{W}_3$, like those with LECs $\bar{\beta}_2$ and $\bar{\beta}_3$ in Eq. (109) and $\bar{\xi}_i$ in Eq. (111), can be absorbed in a redefinition of the coefficients, whose scaling is still given by Eqs. (110) and (112). For operators that transform as $\tilde{V}_4$, the scaling of the coefficients is modified, and they get a contribution from the isospin-symmetry-breaking Lagrangian at order $\Delta = 1, 2$ (listed, for example, in Ref. [32]). Schematically,

$$\bar{\beta}_1 \to \bar{\beta}_1' = \mathcal{O}\left(\left(\tilde{\delta}_0 + \varepsilon \tilde{\delta}_3\right) \frac{m_\pi^2}{M_{\not T}^2}\right), \qquad \bar{\zeta}_i \to \bar{\zeta}_i' = \mathcal{O}\left(\left(\tilde{\delta}_0 + \varepsilon \tilde{\delta}_3\right) \frac{m_\pi^2}{M_{\not T}^2 M_{\rm QCD}}\right). \tag{114}$$

Still at order $\Delta_6 = 1$, interactions arise from the combined effect of the qCEDM and the QCD mass terms. The resulting operators transform as $(\bar{m} S_4 - \varepsilon \bar{m} P_3) \otimes (-\tilde{d}_0 \tilde{V}_4 + \tilde{d}_3 \tilde{W}_3)$. They generate

$$\begin{aligned} \mathcal{L}^{(1)}_{\tilde{q}, f=2} \;=\; & -\frac{\delta \bar{g}_0}{F_\pi D}\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2}\right) \bar{N} \boldsymbol{\tau} \cdot \boldsymbol{\pi} N - \frac{\delta \bar{g}_1}{F_\pi D}\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2}\right) \pi_3 \bar{N} N \\ & - \frac{\bar{g}_2}{F_\pi D} \pi_3 \bar{N} \left(\tau_3 - \frac{2\pi_3}{F_\pi D} \boldsymbol{\tau} \cdot \boldsymbol{\pi}\right) N. \end{aligned} \tag{115}$$

The first two terms are corrections to the isoscalar and isovector $\not P \not T$ couplings in Eq. (105), from which they differ only by terms with three or more pions. The corrections scale as

$$\delta \bar{g}_0 = \mathcal{O}\left(\tilde{\delta}_0 \frac{m_\pi^4}{M_{\not T}^2 M_{\rm QCD}}\right), \qquad \delta \bar{g}_1 = \mathcal{O}\left(\left(\varepsilon \tilde{\delta}_0 + \tilde{\delta}_3\right) \frac{m_\pi^4}{M_{\not T}^2 M_{\rm QCD}}\right). \tag{116}$$

For most practical purposes $\delta \bar{g}_0$ and $\delta \bar{g}_1$ can be absorbed into $\bar{g}_0$ and $\bar{g}_1$. The third operator is the most interesting, as it is the first contribution of the qCEDM to the isospin-breaking pion-nucleon interaction $\pi_3 \bar{N} \tau_3 N$. It has the scaling

$$\bar{g}_2 = \mathcal{O}\left(\varepsilon \tilde{\delta}_3 \frac{m_\pi^4}{M_{\not T}^2 M_{\rm QCD}}\right). \tag{117}$$

Thus, two orders above lowest all three possible non-derivative pion-nucleon $\not P \not T$ interactions receive a contribution from the isovector qCEDM. Just like the $\bar{\theta}$ term [32], the isoscalar qCEDM generates $\bar{g}_2$ as well, but this requires a photon exchange so that $\bar{g}_2$ is suppressed by $\alpha_{\rm em}/4\pi$ and enters three orders above lowest. This provides the second difference between isovector qCEDM and $\bar{\theta}$ term (and isoscalar qCEDM). However, since for all sources $\bar{g}_2$ enters in the subleading Lagrangian, this difference is of little phenomenological interest.

Tadpole elimination affects subleading non-derivative couplings in two ways. First, the elimination of subleading tadpoles shifts the coefficients $\bar{g}_0$ and $\bar{g}_1$ in Eqs. (107) and (108), $\bar{g}_{0,1} \to \bar{g}_{0,1} + \delta' \bar{g}_{0,1}$, with

$$\delta' \bar{g}_0 = \mathcal{O}\left(\left(\tilde{\delta}_0 + \varepsilon \tilde{\delta}_3\right)(1+\varepsilon^2)\frac{m_\pi^4}{M_{\not{T}}^2 M_{\mathrm{QCD}}}\right), \quad \delta' \bar{g}_1 = \mathcal{O}\left(\left(\varepsilon \tilde{\delta}_0 + (1+\varepsilon^2)\tilde{\delta}_3\right)\frac{m_\pi^4}{M_{\not{T}}^2 M_{\mathrm{QCD}}}\right) \tag{118}$$

contributing to the $\Delta_6 = 1$ Lagrangian. Second, the operators in Eq. (115) receive corrections from the transformation of CP-even, chiral-breaking operators that contribute to the chiral Lagrangian at order $\Delta = 3$. These operators are proportional to two powers of the quark masses, and transform as the tensor products $(\bar{m}S_4 - \bar{m}\varepsilon P_3) \otimes (\bar{m}S_4 - \bar{m}\varepsilon P_3)$. The rotation needed to eliminate the leading tapdoles causes them to generate corrections to the coefficients $\delta\bar{g}_{0,1}$ and $\bar{g}_2$. Now $\delta\bar{g}_0$ also depends on $\varepsilon\tilde{\delta}_3$ and $\delta\bar{g}_1$ receives a $\varepsilon^2\tilde{\delta}_3$ correction,

$$\delta\bar{g}_0 \to \delta\bar{g}_0' = \mathcal{O}\left(\left(\tilde{\delta}_0 + \varepsilon \tilde{\delta}_3\right)\frac{m_\pi^4}{M_{\not{T}}^2 M_{\mathrm{QCD}}}\right), \quad \delta\bar{g}_1 \to \delta\bar{g}_1' = \mathcal{O}\left(\left(\varepsilon \tilde{\delta}_0 + (1+\varepsilon^2)\tilde{\delta}_3\right)\frac{m_\pi^4}{M_{\not{T}}^2 M_{\mathrm{QCD}}}\right), \tag{119}$$

while the scaling of $\bar{g}_2$ is unchanged.

6.3 FQLR

The FQLR operator also can generate non-derivative pion-nucleon couplings. Pion-nucleon operators start at $\Delta_6 = -3$, and at this order there is only one operator with the structure of X_{34}, the isovector pion-nucleon interaction. It appears as if the FQLR generates only $\bar{g}_1$ at LO, in stark contrast with the $\bar{\theta}$ term and qCEDM which generate, respectively, $\bar{g}_0$ and $\bar{g}_{0,1}$ at LO. However, the removal of the tadpole in Eq. (78) effectively causes the appearance of the $\bar{g}_0$ interaction at $\Delta_6 = -3$. At this order,

$$\mathcal{L}^{(-3)}_{\mathrm{LR},f=2} = -\frac{\bar{g}_0}{F_\pi D}\bar{N}\boldsymbol{\tau}\cdot\boldsymbol{\pi}N - \frac{\bar{g}_1}{F_\pi D}\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right)\pi_3\bar{N}N - \frac{\bar{g}_1'}{F_\pi D}\pi_3\bar{N}N, \tag{120}$$

where

$$\bar{g}_1 = \mathcal{O}\left(\xi\frac{M_{\mathrm{QCD}}^3}{M_{\not{T}}^2}\right) \tag{121}$$

arises directly, and

$$\bar{g}_0 = \delta m_N \frac{\bar{\Delta}^{(-4)}_{\mathrm{LR}}}{m_\pi^2} = \mathcal{O}\left(\varepsilon\xi\frac{M_{\mathrm{QCD}}^3}{M_{\not{T}}^2}\right), \qquad \bar{g}_1' = 2\Delta m_N \frac{\bar{\Delta}^{(-4)}_{\mathrm{LR}}}{m_\pi^2} = \mathcal{O}\left(\xi\frac{M_{\mathrm{QCD}}^3}{M_{\not{T}}^2}\right) \tag{122}$$

stem from tadpole removal via Eq. (104). One can eliminate $\bar{\Delta}^{(-4)}_{\mathrm{LR}}$ and write

$$\bar{g}_0 = \frac{\delta m_N}{2\Delta m_N}\bar{g}_1'. \tag{123}$$

A lattice evaluation found $\delta m_N = 2.26$ MeV [60], while values for Δm_N range between 45 and 60 MeV [61]. Therefore, for FQLR $\bar{g}_0$ is actually only a few percent of $\bar{g}_1'$.

At higher orders FQLR generates some interactions that resemble those of the isovector qCEDM, although with a more complicated structure, plus some new interactions. At $\Delta_6 = -2$ and $\Delta_6 = -1$ we find interactions similar to those with LECs $\bar{\beta}_{2,3}$ in Eq. (109) and $\bar{\xi}_i$ in Eq. (111), but with the replacements

$$\begin{aligned} \frac{2\pi_3}{F_\pi D} &\to \frac{2\pi_3}{F_\pi D}\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right) \\ \left(\delta_{i3} - \frac{2\pi_3\pi_i}{F_\pi^2 D}\right) &\to \left[\delta_{i3}\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right) - \frac{2\pi_i\pi_3}{F_\pi^2 D}\left(3 - \frac{4\boldsymbol{\pi}^2}{F_\pi^2 D}\right)\right]. \end{aligned} \tag{124}$$

The additional interactions are

$$\mathcal{L}^{(-2)}_{\mathrm{LR},f=2} = \frac{2\bar{\beta}_4}{F_\pi^2 D}\left[(D_\mu\pi_3)\pi_i + (\boldsymbol{\pi}\cdot D_\mu\boldsymbol{\pi})\left(\delta_{i3} - \frac{4\pi_i\pi_3}{F_\pi^2 D}\right)\right]\bar{N}S^\mu\tau_i N, \tag{125}$$

and

$$\begin{aligned} \mathcal{L}^{(-1)}_{\mathrm{LR},f=2} = {} & \frac{2\bar{\xi}_{12}}{F_\pi^3 D}\left[D^\mu\pi_3 - \frac{2\pi_3}{F_\pi^2 D}(\boldsymbol{\pi}\cdot D_\mu\boldsymbol{\pi})\right](\boldsymbol{\pi}\cdot D_\mu\boldsymbol{\pi})\bar{N}N \\ & + \frac{2\bar{\xi}_{13}}{F_\pi^3 D}\left[\delta_{i3}\pi_j + \delta_{j3}\pi_i - \frac{4\pi_3\pi_i\pi_j}{F_\pi^2 D}\right](D_\mu\boldsymbol{\pi}\times D_\nu\boldsymbol{\pi})_i\,\bar{N}i[S^\mu, S^\nu]\tau_j N \\ & + \frac{2i\bar{\xi}_{14}}{F_\pi^2 D}\left[\delta_{i3}\pi_j + \delta_{j3}\pi_i - \frac{4\pi_3\pi_i\pi_j}{F_\pi^2 D}\right](v\cdot D\pi_i)\bar{N}\tau_j S\cdot\mathcal{D}_- N, \end{aligned} \tag{126}$$

where

$$\bar{\beta}_{2,3,4} = \mathcal{O}\left(\xi\frac{M_{\mathrm{QCD}}^2}{M_{\not{T}}^2}\right), \qquad \bar{\xi}_i = \mathcal{O}\left(\xi\frac{M_{\mathrm{QCD}}}{M_{\not{T}}^2}\right). \tag{127}$$

The RPI relations are the same as in Eq. (113), with the additional relation

$$\bar{\xi}_{14} = -\frac{\bar{\beta}_4}{m_N}. \tag{128}$$

Just like for qCEDM, tadpole removal induces the subleading isospin-conserving $\not{P}\not{T}$ operators with LECs $\bar{\beta}_1$ in Eq. (109) and $\bar{\zeta}_i$ in Eq. (111), but with scalings

$$\bar{\beta}_1 = \mathcal{O}\left(\varepsilon\xi\frac{M_{\mathrm{QCD}}^2}{M_{\not{T}}^2}\right), \qquad \bar{\zeta}_i = \mathcal{O}\left(\varepsilon\xi\frac{M_{\mathrm{QCD}}}{M_{\not{T}}^2}\right). \tag{129}$$

Again, two orders above the lowest order, operators appear due to an insertion of the quark mass (difference),

$$\begin{aligned} \mathcal{L}^{(-1)}_{\mathrm{LR},f=2} = {} & -\frac{\delta\bar{g}_0}{F_\pi D}\bar{N}\boldsymbol{\tau}\cdot\boldsymbol{\pi}N - \frac{1}{F_\pi D}\left[\delta_1\bar{g}_1 + \delta_2\bar{g}_1\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right)^2\right]\pi_3\bar{N}N \\ & -\frac{\bar{g}_2}{F_\pi D}\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right)\pi_3\bar{N}\left(\tau_3 - \frac{2\pi_3}{F_\pi D}\boldsymbol{\tau}\cdot\boldsymbol{\pi}\right)N - \frac{\bar{g}_3}{F_\pi^3 D^3}\pi_3^2\bar{N}\boldsymbol{\tau}\cdot\boldsymbol{\pi}N, \end{aligned} \tag{130}$$

where the LECs scale as

$$\delta_{1,2}\bar{g}_1 = \mathcal{O}\left(\xi\frac{m_\pi^2 M_{\rm QCD}}{M_{\not T}^2}\right), \qquad \delta\bar{g}_0 = \mathcal{O}\left(\varepsilon\xi\frac{m_\pi^2 M_{\rm QCD}}{M_{\not T}^2}\right), \qquad \bar{g}_{2,3} = \mathcal{O}\left(\varepsilon\xi\frac{m_\pi^2 M_{\rm QCD}}{M_{\not T}^2}\right). \tag{131}$$

Although the operators in Eq. (130) appear to be very complicated, if we ignore terms with three or more pions the $\delta_{1,2}\bar{g}_1$ and $\delta\bar{g}_0$ couplings are contributions to the standard non-derivative pion-nucleon interactions. Just as for the isovector qCEDM, the third pion-nucleon coupling $\bar{g}_2$ comes in two orders higher than $\bar{g}_{0,1}$.

As in the case of the qCEDM, the elimination of the tadpoles modifies Eq. (130). The elimination of the subleading tadpoles $\bar{\Delta}^{(-2)}_{\rm LR1}$ and $\bar{\Delta}^{(-2)}_{\rm LR2}$ shifts $\delta\bar{g}_0$ and $\delta_1\bar{g}_1$, but does not modify their dependence on ξ and ε. On the other hand, the effect of the elimination of the leading tadpole on chiral-breaking operators proportional to two powers of the quark masses $\bar{m}$ and $\bar{m}\,\varepsilon$ is to generate a contribution identical to Eq. (115), with the coefficients replaced by

$$\delta'\bar{g}_1 = \mathcal{O}\left(\left(1+\varepsilon^2\right)\xi\frac{m_\pi^2 M_{\rm QCD}}{M_{\not T}^2}\right), \qquad \delta'\bar{g}_0 = \mathcal{O}\left(\varepsilon\xi\frac{m_\pi^2 M_{\rm QCD}}{M_{\not T}^2}\right), \qquad \bar{g}_2' = \mathcal{O}\left(\varepsilon\xi\frac{m_\pi^2 M_{\rm QCD}}{M_{\not T}^2}\right). \tag{132}$$

6.4 qEDM

The pion-nucleon interactions originating from the qEDM arise from the tensor product $(-d_0 V_4 + d_3 W_3)\otimes e(I^\mu/6 + T^\mu_{34}/2)$, and thus have a rich chiral structure:

$$\mathcal{L}^{(2)}_{q,f=2} = -\frac{\bar{g}_0}{F_\pi D}\bar{N}\boldsymbol{\tau}\cdot\boldsymbol{\pi}N - \frac{\bar{g}_1}{F_\pi D}\pi_3\bar{N}N - \frac{\bar{g}_2}{F_\pi D}\pi_3\bar{N}\left[\tau_3 + \frac{2}{F_\pi^2 D}\left(\pi_3\boldsymbol{\tau}\cdot\boldsymbol{\pi} - \boldsymbol{\pi}^2\tau_3\right)\right]N, \tag{133}$$

the scaling of the LECs being

$$\begin{aligned}\bar{g}_0 &= \mathcal{O}\left(\left(\delta_0+\delta_3\right)\frac{\alpha_{\rm em}}{4\pi}\frac{m_\pi^2 M_{\rm QCD}}{M_{\not T}^2}\right), \qquad \bar{g}_1 = \mathcal{O}\left(\left(\delta_0+\delta_3\right)\frac{\alpha_{\rm em}}{4\pi}\frac{m_\pi^2 M_{\rm QCD}}{M_{\not T}^2}\right),\\ \bar{g}_2 &= \mathcal{O}\left(\delta_3\frac{\alpha_{\rm em}}{4\pi}\frac{m_\pi^2 M_{\rm QCD}}{M_{\not T}^2}\right).\end{aligned} \tag{134}$$

The qEDM also generates tadpoles, as given in Eq. (82), which can be removed in the same way as for the other sources. This removal alters the ε dependence of $\bar{g}_0$, in a way that can be summarized by replacing $(\delta_0+\delta_3)$ with $(\delta_0+\delta_3)(1+\varepsilon)$ in Eq. (134). Since, as Eq. (134) shows, $\bar{g}_{0,1}$ already receive contributions from both the isoscalar and isovector qEDMs, the added contributions are not particularly relevant.

Note that all possible non-derivative pion-nucleon interactions appear at the same order. (This only holds for the isovector qEDM, since for the isoscalar qEDM $\bar{g}_2$ is suppressed.) Only the qEDM has this property: for all other $\not P\not T$ dimension-six sources $\bar{g}_2$ is suppressed with respect to $\bar{g}_0$ and/or $\bar{g}_1$. However, the $\alpha_{\rm em}/4\pi$ suppression makes all the qEDM pion-nucleon couplings pretty unimportant.

6.5 χISs

In the pion-nucleon sector, no chiral-invariant $\not{P}\not{T}$ operator can be constructed with zero or one covariant derivative. The first operators therefore start at $\Delta_6 = -1$ and have two covariant derivatives or one insertion of the quark mass (difference):

$$\begin{aligned}\mathcal{L}^{(-1)}_{w,f=2} &= \frac{\bar{\zeta}_1}{F_\pi}(\mathcal{D}_{\mu,\perp} D^\mu_\perp \boldsymbol{\pi}) \cdot \bar{N}\boldsymbol{\tau}N + \frac{\bar{\zeta}_2}{F_\pi^2}(D_\mu\boldsymbol{\pi} \times v \cdot D\boldsymbol{\pi}) \cdot \bar{N}S^\mu\boldsymbol{\tau}N \\ &\quad - \frac{\bar{g}_0}{F_\pi D}\bar{N}\boldsymbol{\tau}\cdot\boldsymbol{\pi}N - \frac{\bar{g}_1}{F_\pi D}\pi_3\bar{N}N, \end{aligned} \tag{135}$$

with scalings

$$\bar{\zeta}_i = \mathcal{O}\left(w\frac{M_{\mathrm{QCD}}}{M^2_{\not{T}}}\right), \qquad \bar{g}_0 = \mathcal{O}\left(w\frac{m^2_\pi M_{\mathrm{QCD}}}{M^2_{\not{T}}}\right), \qquad \bar{g}_1 = \mathcal{O}\left(\varepsilon w\frac{m^2_\pi M_{\mathrm{QCD}}}{M^2_{\not{T}}}\right). \tag{136}$$

It is interesting to notice that rotation invariance alone would allow a two-derivative operator of the form $\bar{\zeta}_4(D_\nu\boldsymbol{\pi}) \cdot \bar{N}\boldsymbol{\tau}[S^\mu, S^\nu]\mathcal{D}_{\mu-}N$ in Eq. (135). However, this operator is not RPI by itself, and its variation cannot be absorbed by any other operator in the leading-order Lagrangian. RPI, therefore, forces $\bar{\zeta}_4$ to vanish in leading order. The operator $\bar{\zeta}_4$ appears at NNLO, and it is linked to $\bar{g}_0$ by RPI, as in Eq. (113).

Here the tadpole rotation induces the shifts

$$\begin{aligned}\bar{g}_0 &\to \bar{g}_0 + \delta m_N\frac{\bar{\Delta}^{(-2)}_w}{m^2_\pi} = \mathcal{O}\left(\left(1+\varepsilon^2\right)w\frac{m^2_\pi M_{\mathrm{QCD}}}{M^2_{\not{T}}}\right), \qquad (137)\\ \bar{g}_1 &\to \bar{g}_1 + \Delta m_N\frac{\bar{\Delta}^{(-2)}_w}{m^2_\pi} = \mathcal{O}\left(\varepsilon w\frac{m^2_\pi M_{\mathrm{QCD}}}{M^2_{\not{T}}}\right), \qquad (138)\end{aligned}$$

which are less interesting, because the dependence on w is not fundamentally changed. As before, we absorb contributions from tadpole elimination in $\bar{g}_i$

From Eq. (135) we see that the χISs, just as the qCEDM and in contrast to the $\bar{\theta}$ term, induce $\bar{g}_1$ at the same order as $\bar{g}_0$. Differently from all the other sources, the χISs also generate two-derivative operators of the same importance as $\bar{g}_0$ and $\bar{g}_1$.

One order higher we find various types of interactions: chiral-invariant operators with three covariant derivatives; chiral-symmetry-breaking operators with one power of quark masses and one covariant derivative; and electromagnetic operators coming from the tensor product $d_W\bar{I} \otimes e(I^\mu/6 + T^\mu_{34}/2) \otimes e(I^\mu/6 + T^\mu_{34}/2)$. The chiral-invariant operators are

$$\begin{aligned}\mathcal{L}^{(0)}_{w,f=2} &= -\frac{i\bar{\varkappa}_1}{F_\pi}\mathcal{D}_\mu(v\cdot D\boldsymbol{\pi}) \cdot \bar{N}\boldsymbol{\tau}\mathcal{D}^\mu_{\perp-}N + \frac{i\varkappa_2}{2F_\pi^2}(D_\mu\boldsymbol{\pi} \times D_\nu\boldsymbol{\pi}) \cdot \bar{N}\boldsymbol{\tau}S^\mu\mathcal{D}^\nu_{\perp-}N \\ &\quad + \frac{\bar{\varkappa}_3}{F_\pi^2}\left(\mathcal{D}_\mu(D\boldsymbol{\pi})^2\right)\bar{N}S^\mu N + \frac{\bar{\varkappa}_4}{F_\pi^2}\left(\mathcal{D}^\mu_\perp(D_\mu\boldsymbol{\pi}\cdot D_\nu\boldsymbol{\pi})\right)\bar{N}S^\nu N, \end{aligned} \tag{139}$$

where

$$\bar{\varkappa}_{3,4} = \mathcal{O}\left(\frac{w}{M^2_{\not{T}}}\right), \tag{140}$$

while RPI gives

$$\bar{\varkappa}_{1,2} = \frac{\bar{\zeta}_{1,2}}{m_N}. \tag{141}$$

The quark-mass insertions give rise to

$$\begin{aligned}\mathcal{L}^{(0)}_{w,f=2} &= \frac{2\bar{\beta}_1}{F_\pi^2 D}(\boldsymbol{\pi}\cdot\mathcal{D}_\mu\boldsymbol{\pi})\bar{N}S^\mu N + \frac{2\bar{\beta}_2}{F_\pi^2 D}(\pi_3\mathcal{D}_\mu\boldsymbol{\pi})\cdot\bar{N}S^\mu\boldsymbol{\tau}N \\ &\quad + \frac{\bar{\beta}_3}{F_\pi}\left(\delta_{i3} - \frac{2\pi_i\pi_3}{F_\pi^2 D}\right)\bar{N}(\boldsymbol{\tau}\times v\cdot D\boldsymbol{\pi})_i N, \end{aligned} \tag{142}$$

with

$$\bar{\beta}_1 = \mathcal{O}\left(w\frac{m_\pi^2}{M_{\not{T}}^2}\right), \qquad \bar{\beta}_{2,3} = \mathcal{O}\left(\varepsilon w\frac{m_\pi^2}{M_{\not{T}}^2}\right). \tag{143}$$

The removal of tadpoles does not affect the chiral-invariant operators in Eq. (139), while it slightly modifies the ε dependence of β_1 in Eq. (142), in a way analogous to Eq. (137).

Finally, integrating out a hard photon gives the single operator

$$\mathcal{L}^{(0)}_{w,f=2} = -\frac{\bar{g}_2}{F_\pi D}\bar{N}\left\{\pi_3\tau_3 - \boldsymbol{\tau}\cdot\boldsymbol{\pi}\left[1 - \frac{2}{F_\pi^2 D}\left(\boldsymbol{\pi}^2 - \pi_3^2\right)\right]\right\}N. \tag{144}$$

We see that at this order the first contribution to the non-derivative pion-nucleon interaction $\bar{g}_2$ appears. It scales as

$$\bar{g}_2 = \mathcal{O}\left(w\frac{\alpha_{\text{em}}}{4\pi}\frac{M_{\text{QCD}}^3}{M_{\not{T}}^2}\right). \tag{145}$$

With the usual counting $\alpha_{\text{em}}/4\pi \sim m_\pi^3/M_{\text{QCD}}^3$, this interaction is suppressed by one order compared to the $\bar{g}_{0,1}$ interactions.

7 The pion-nucleon form factor

An important element in the evaluation of hadronic and nuclear EDMs is the $\not{P}\not{T}$ pion-nucleon coupling. In this section, we summarize the pion-nucleon interactions for the different $\not{P}\not{T}$ sources we have considered, by calculating the $\not{P}\not{T}$ PNFF with the Lagrangian derived above. For convenience, we consider the Lagrangian after tadpole extermination.

We consider the three-point Green's function for an incoming (outgoing) nucleon of momentum p^μ (p'^μ) and an outgoing pion of momentum $q^\mu = p^\mu - p'^\mu$ and isospin a. We take the incoming and outgoing nucleon to be nonrelativistic and on-shell, so

$$p^0 = \frac{\vec{p}^2}{2m_N} - \Delta m_N \mp \frac{\delta m_N}{2} + \ldots, \quad p'^0 = \frac{\vec{p}'^2}{2m_N} - \Delta m_N \mp \frac{\delta m_N}{2} + \ldots, \tag{146}$$

where the $-(+)$ signs holds for protons (neutrons) and the arrow denotes vectors in three-dimensional Euclidean space, $p^\mu = (p^0, \vec{p})$, $p_\mu = (p^0, -\vec{p})$. It is clear that p^0 enters at one order higher than $\vec{p}$. The Green function for on-shell nucleons can be parameterized by three form factors, corresponding to three different isospin structures,

$$V_a(q,K) = -\frac{i}{F_\pi}\left[F_1(q,K)\tau_a + F_2(q,K)\delta_{a3} + F_3(q,K)\delta_{a3}\tau_3\right], \tag{147}$$

in terms of the functions $F_{1,2,3}$ of q^μ and $K^\mu = (p^\mu + p'^\mu)/2$. In what follows, we give F_1, F_2, and F_3 for each of the dimension-six $\not{P}\not{T}$ sources. We calculate the PNFFs up to the order where each of them gets a non-vanishing contribution. As derived in Sec. 6, this implies going to NNLO for qCEDM and FQLR, and to NLO for χISs. For qEDM we need the LO contributions only.

At LO and NLO, for all sources but FQLR contributions arise exclusively from tree diagrams. At NLO for the FQLR and at NNLO for the other sources loops appear. We use dimensional regularization in d spacetime dimensions, which introduces the renormalized scale μ and

$$L = \frac{2}{4-d} - \gamma_E + \ln 4\pi, \tag{148}$$

where $\gamma_E \simeq 0.557$ is the Euler-Mascheroni constant. At this order we use $q^0 = 0$.

7.1 qEDM

The PNFFs from the qEDM are very simple to the order we are interested in, since all three PNFFs appear at the same, leading order: $\Delta_6 = 2$. We read off from the Lagrangian directly,

$$F_1 = \bar{g}_0, \qquad F_2 = \bar{g}_1, \qquad F_3 = \bar{g}_2. \tag{149}$$

Since these interactions contain suppression factors of $\alpha_{\rm em}/4\pi \sim m_\pi^3/M_{\rm QCD}^3$, they are most likely not important in calculations of nuclear EDMs, which are dominated by short-range contributions to the nucleon EDM [36].

7.2 χISs

For the gCEDM and chiral-invariant four-quark operators, the PNFFs receive contributions at LO and NLO from interactions with LECs, respectively, $\bar{g}_0$, $\bar{g}_1$, and $\bar{\zeta}_1$ in Eq. (135), and $\bar{g}_2$ in Eq. (144). The contributions to the PNFFs can be read off easily,

$$\begin{aligned} F_1(\vec{q}) &= \bar{g}_0 - \bar{g}_2 - \bar{\zeta}_1 \vec{q}^{\,2}, & (150)\\ F_2 &= \bar{g}_1, & (151)\\ F_3 &= \bar{g}_2. & (152)\end{aligned}$$

We conclude that, for the chiral-invariant $\not{P}\not{T}$ sources, F_1 and F_2 appear at the same order and F_3 appears one order down in the $Q/M_{\rm QCD}$ expansion. Apart from contributing to F_3, $\bar{g}_2$ also contributes to F_1. This is of no phenomenological interest, since F_1 receives a larger contribution from $\bar{g}_0$. At LO F_1 depends on the pion momentum because of the presence of the chiral-invariant operator $\bar{\zeta}_1$. The consequences for the $\not{P}\not{T}$ NN potential have been worked out in Ref. [35], where it was shown that $\bar{\zeta}_1$ generates a long-range $\not{P}\not{T}$ potential, which can be accounted for by a redefinition of $\bar{g}_0$, and a short-range $\not{P}\not{T}$ potential, which can be absorbed in a nucleon-nucleon contact interaction.

7.3 qCEDM

In LO ($\Delta_6 = -1$) and NLO ($\Delta_6 = 0$) the PNFF arises from tree diagrams where the pion-nucleon vertices are, respectively, the two non-derivative interactions with LECs $\bar{g}_0$ and $\bar{g}_1$ in



Figure 1: One-loop contributions of relative order $\mathcal{O}(m_\pi^2/(2\pi F_\pi)^2)$ to the pion-nucleon form factors $F_1(\vec{q},\vec{K})$ and $F_2(\vec{q},\vec{K})$. A nucleon (pion) is represented by a solid (dashed) line; the $\not{P}\not{T}$ vertices from Eq. (105) are indicated by a square. The other vertices represent LO interactions from Eq. (63). For simplicity only one possible ordering is shown.

Eq. (105), and the single-derivative interaction with LEC $\bar{\beta}_3$ in Eq. (109). At NNLO ($\Delta_6 = 1$) there are further tree-level contributions from the two-derivative interactions with LECs $\bar{\zeta}_1$, $\bar{\zeta}_4$, $\bar{\zeta}_6$, $\bar{\xi}_1$, $\bar{\xi}_4$, $\bar{\xi}_5$ and $\bar{\xi}_7$ in Eq. (111), and the non-derivative interactions with LECs $\delta\bar{g}_0$, $\delta\bar{g}_1$ and $\bar{g}_2$ in Eq. (115). Use of the RPI relations (113) shows that the PNFFs can be expressed in terms of $\bar{g}_0$, $\bar{g}_1$, $\bar{\beta}_3$, $\bar{\zeta}_1$, $\bar{\xi}_1$, $\delta\bar{g}_0$, $\delta\bar{g}_1$ and $\bar{g}_2$. The operators with LECs $\bar{g}_0$, $\bar{\zeta}_1$ and $\delta\bar{g}_0$ contribute to F_1, those with LECs $\bar{g}_1$, $\bar{\xi}_1$, $\delta\bar{g}_1$ to F_2, and that with LEC $\bar{g}_2$ to F_3. The operator with LEC $\bar{\beta}_3$ and its recoil correction contribute to the PNFF

$$V_a(\vec{q},\vec{K}) = \frac{\bar{\beta}_3}{F_\pi}\epsilon_{3ab}\tau_b\left(q^0 - \frac{\vec{K}\cdot\vec{q}}{m_N}\right). \tag{153}$$

When we use the nucleon on-shell conditions (146), this contributes to F_1 and F_3.

At NNLO there are also one-loop diagrams where the $\not{P}\not{T}$ vertex is one of the two non-derivative interactions with LECs $\bar{g}_0$ and $\bar{g}_1$ in Eq. (105), other vertices coming from the LO chiral Lagrangian, Eq. (63). These loops are shown in Fig. 1. The structure of the diagrams is such that the momentum of the external pion never flows into the loop and the only scale in the integral is the pion mass. As a consequence, the diagrams do not yield any non-trivial momentum dependence and simply renormalize the LECs $\delta\bar{g}'_{0,1}$ and $\delta'\bar{g}_{0,1}$. We define the renormalized LECs

$$\begin{aligned}
\bar{\delta}\bar{g}_0 &= \delta\bar{g}'_0 + \delta'\bar{g}_0 + \frac{\bar{g}_0}{4}\frac{m_\pi^2}{(2\pi F_\pi)^2}\left[(1+3g_A^2)\left(L+1-\log\frac{m_\pi^2}{\mu^2}\right) - 2g_A^2\right], \\
\bar{\delta}\bar{g}_1 &= \delta\bar{g}'_1 + \delta'\bar{g}_1 + \frac{\bar{g}_1}{4}\frac{m_\pi^2}{(2\pi F_\pi)^2}\left[(5-9g_A^2)\left(L+1-\log\frac{m_\pi^2}{\mu^2}\right) + 6g_A^2\right].
\end{aligned} \tag{154}$$

With this definition the PNFFs for on-shell nucleons read, up to NNLO,

$$F_1(\vec{q},\vec{K}) = \bar{g}_0\left\{1 - \frac{1}{2m_N^2}\left[\vec{K}^{\,2} + \vec{S}\cdot\left(\vec{K}\times\vec{q}\right)\right]\right\} + \bar{\delta}\bar{g}_0 + \delta m_N\bar{\beta}_3 - \bar{\zeta}_1\vec{q}^{\,2}, \tag{155}$$

$$F_2(\vec{q},\vec{K}) = \bar{g}_1\left\{1 - \frac{1}{2m_N^2}\left[\vec{K}^{\,2} + \vec{S}\cdot\left(\vec{K}\times\vec{q}\right)\right]\right\} + \bar{\delta}\bar{g}_1 - \bar{\xi}_1\vec{q}^{\,2}, \tag{156}$$

$$F_3 = \bar{g}_2 - \delta m_N\bar{\beta}_3. \tag{157}$$

We see that F_1 and F_2 receive contributions at the same order. Two orders down we find momentum dependence of these PNFFs, as well as the first static contribution to F_3. The implication of the momentum dependence to the two-nucleon potential was discussed in Ref. [35].



Figure 2: One-loop contributions from purely mesonic $\not{P}\not{T}$ interactions to the pion-nucleon form factors. The square denotes the $\not{P}\not{T}$ vertex from Eq. (100). The other notation is as in Fig. 1.

7.4 FQLR

The main part of the PNFFs from FQLR is very similar to those from qCEDM. After replacing the scalings of the LECs, one sees that contributions to F_1 and F_2 start at LO ($\Delta_6 = -3$), and contributions to F_3 at NNLO ($\Delta_6 = -1$); at this order F_1 and F_2 obtain analytic momentum dependence.

Again, the diagrams in Fig. 1 only renormalize LECs. The different chiral structure of the isovector vertices $\bar{g}_1$ and $\bar{g}_1'$ in Eq. (120) only affects the first diagram in Fig. 1, with the minor consequence of modifying the counterterm $\bar{\delta}\bar{g}_1$ with respect to Eq. (154). The results in Eqs. (155), (156), and (157) give the dominant contributions to the three PNFFs, with the replacement $\bar{g}_1 \to \bar{g}_1 + \bar{g}_1'$. and

$$\begin{aligned}\bar{\delta}\bar{g}_1 &= \delta_1\bar{g}_1 + \delta_2\bar{g}_1 + \delta'\bar{g}_1 + \frac{5}{4}\left(3\bar{g}_1 + \bar{g}_1'\right)\frac{m_\pi^2}{(2\pi F_\pi)^2}\left(L + 1 - \log\frac{m_\pi^2}{\mu^2}\right) \\ &\quad + \frac{3g_A^2}{4}\left(\bar{g}_1 + \bar{g}_1'\right)\frac{m_\pi^2}{(2\pi F_\pi)^2}\left[-3\left(L + 1 - \log\frac{m_\pi^2}{\mu^2}\right) + 2\right]. \end{aligned} \tag{158}$$

However, this is not the whole story. As shown in Sec. 5.5, the elimination of the tadpoles in case of the FQLR leaves the three-pion vertex with LEC $\bar{\Delta}_{\mathrm{LR}}^{(-4)}$ in Eq. (100) with a lower chiral index than the dominant pion-nucleon interactions. The one-loop diagrams in Fig. 2 contribute to the PNFFs at NLO ($\Delta_6 = -2$) and add to F_2 a non-analytic momentum dependence not present for the qCEDM. This PNFF becomes, instead of Eq. (156),

$$\begin{aligned} F_2(\vec{q}, \vec{K}) &= \left(\bar{g}_1 + \bar{g}_1'\right)\left\{1 - \frac{1}{2m_N^2}\left[\vec{K}^{\,2} + \vec{S}\cdot\left(\vec{K}\times\vec{q}\right)\right]\right\} + \bar{\delta}\bar{g}_1 - \bar{\xi}_1\vec{q}^{\,2} \\ &\quad - 5\pi g_A^2\frac{m_\pi\bar{\Delta}_{\mathrm{LR}}^{(-4)}}{(2\pi F_\pi)^2}\, f_2\left(\frac{|\vec{q}|}{2m_\pi}\right), \end{aligned} \tag{159}$$

where

$$f_2(x) = 1 + \frac{1 + 2x^2}{2x}\arctan x. \tag{160}$$

Using Eq. (122) we can eliminate $\Delta_{\mathrm{LR}}^{(-4)}$ in favor of $\bar{g}_1'$ or $\bar{g}_0$. Precise measurements on the deuteron $\not{P}\not{T}$ electromagnetic form factors, which depend strongly on F_2 [59, 62, 34], could, in principle, measure this momentum dependence and separate the FQLR from the qCEDM.

8 Electromagnetic interactions

The experimental interest in EDMs bring $\not{P}\not{T}$ electromagnetic interactions to the forefront. Some interactions of hadrons with soft photons can be obtained using the $U(1)$-gauge covariant derivatives (68) in existing operators. More interesting are the interactions that arise through the field strength $F_{\mu\nu}$, which we describe here.

Since the pion has spin 0, we cannot construct an EDM operator in the $f = 0$ sector. In contrast, there are plenty of $\not{P}\not{T}$ interactions in the $f = 2$ sector.

8.1 qCEDM

In the case of the qCEDM, operators containing the electromagnetic field strength have the chiral properties of the tensor product of the $\not{P}\not{T}$ source and the CP-even electromagnetic interactions. In lowest orders we need only $(-\tilde{d}_0\tilde{V}_4 + \tilde{d}_3\tilde{W}_3) \otimes e(I^\mu/6 + T^\mu_{34}/2)$.

Interactions start at $\Delta_6 = 1$ and transform as the fourth component of a vector, or as the product of a vector and an anti-symmetric tensor,

$$\begin{aligned}
\mathcal{L}^{(1)}_{\tilde{q},f=2\,\mathrm{em}} \;=\;& -2\bar{N}\left[\bar{d}_0\left(1-\frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right)+\bar{d}_1\left(\tau_3-\frac{2\pi_3}{F_\pi^2 D}\boldsymbol{\tau}\cdot\boldsymbol{\pi}\right)\right]S^\mu N\, v^\nu F_{\mu\nu} \\
& -2\bar{d}_1'\left(1-\frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right)\bar{N}\left[\tau_3-\frac{2}{F_\pi^2 D}\left(\boldsymbol{\pi}^2\tau_3-\pi_3\boldsymbol{\pi}\cdot\boldsymbol{\tau}\right)\right]S^\mu N\, v^\nu F_{\mu\nu} \\
& +\frac{1}{F_\pi D}\epsilon^{\mu\nu\alpha\beta}v_\alpha\bar{N}\left(\bar{c}_0\boldsymbol{\tau}\cdot\boldsymbol{\pi}+\bar{c}_1\pi_3\right)S_\beta N\, F_{\mu\nu} \\
& +\bar{c}_2\frac{\pi_3}{F_\pi D}\epsilon^{\mu\nu\alpha\beta}v_\alpha\bar{N}\left[\tau_3-\frac{2}{F_\pi^2 D}\left(\boldsymbol{\pi}^2\tau_3-\pi_3\boldsymbol{\pi}\cdot\boldsymbol{\tau}\right)\right]S_\beta N\, F_{\mu\nu},
\end{aligned} \tag{161}$$

where the electromagnetic LECs scale as

$$\begin{aligned}
\bar{d}_{0,1}, \bar{c}_{0,1} &= \mathcal{O}\left(e\left(\tilde{\delta}_0+\tilde{\delta}_3\right)\frac{m_\pi^2}{M_{\not{T}}^2 M_{\mathrm{QCD}}}\right), \\
\bar{d}_1' = \mathcal{O}\left(e\tilde{\delta}_0\frac{m_\pi^2}{M_{\not{T}}^2 M_{\mathrm{QCD}}}\right), &\qquad \bar{c}_2 = \mathcal{O}\left(e\tilde{\delta}_3\frac{m_\pi^2}{M_{\not{T}}^2 M_{\mathrm{QCD}}}\right).
\end{aligned} \tag{162}$$

The first term in Eq. (161) is a short-range contribution to the isoscalar nucleon EDM. The second and third terms both contribute to the isovector nucleon EDM, and they differ only by interactions involving two or more pions, such that separating them is practically impossible. The last three terms in Eq. (161) are pion-nucleon-photon interactions which play a role in the calculation of the deuteron MQM [34, 37].

As always, the isoscalar qCEDM generates the same interactions as the $\bar{\theta}$ term [32]. But, while in the pion and pion-nucleon sectors the isoscalar qCEDM generated only isoscalar interactions in LO, in the electromagnetic sector both isoscalar and isovector interactions appear, due to breaking of chiral symmetry by the quark electric charge. The main extra feature of the isovector qCEDM is the appearance of the $\bar{c}_2$ interaction, which has a $\tau_3\pi_3$ structure. However, it is unlikely that this operator has any important physical consequences.

The elimination of the leading tadpole modifies to coefficients in Eq. (161) in a way that can be schematically summarized by the replacement $\tilde{\delta}_0 \to \tilde{\delta}_0 + \varepsilon\tilde{\delta}_3$ in Eq. (162). Since most operators already receive contributions from both the isoscalar and isovector qCEDM these shifts are not particularly interesting.

The Schiff moment of the nucleon [43, 33] and EDMs of light nuclei [36] are dominated by $\not{P}\not{T}$ pion-nucleon interactions. The LO $\not{P}\not{T}$ nucleon-photon operators have little impact on these observables, so we do not construct operators with higher chiral index, which would be even more suppressed.

8.2 FQLR

The interactions stemming from FQLR are similar to those from the qCEDM. Already at LO ($\Delta_6 = -1$) FQLR generates operators with a more complicated pion structure due to the $X_{34} \otimes T^{\mu}_{34}/2$ tensor product in $\mathrm{Im}\,\Xi\, X_{34} \otimes e(I^{\mu}/6 + T^{\mu}_{34}/2)$. However, apart from terms with two or more pions, the Lagrangian is identical to Eq. (161):

$$\mathcal{L}^{(-1)}_{\mathrm{LR},f=2\ \mathrm{em}} = \mathcal{L}^{(1)}_{\tilde{q},f=2\ \mathrm{em}} + \dots, \tag{163}$$

but with the scaling

$$\bar{d}_{0,1},\, \bar{d}_1',\, \bar{c}_{0,1},\, \bar{c}_2 = \mathcal{O}\left(e\xi \frac{M_{\mathrm{QCD}}}{M_{\not{T}}^2}\right). \tag{164}$$

Tadpole extermination brings in no important new features. Like for the qCEDM, operators with additional pions and/or higher chiral index have no obvious phenomenological interest.

8.3 qEDM

Since the qEDM contains a photon field, it yields nucleon-photon operators that transform like itself, namely as the third and fourth components of, respectively, the vectors V and W in Eq. (50). At LO this generates

$$\begin{aligned}\mathcal{L}^{(1)}_{q,f=2\ \mathrm{em}} &= -2\bar{N}\left[\bar{d}_0\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right) + \bar{d}_1\left(\tau_3 - \frac{2\pi_3}{F_\pi^2 D}\boldsymbol{\tau}\cdot\boldsymbol{\pi}\right)\right] S^{\mu} N\, v^{\nu} F_{\mu\nu} \\ &\quad + \frac{1}{F_\pi D}\epsilon^{\mu\nu\alpha\beta} v_\alpha \bar{N}(\bar{c}_0 \boldsymbol{\tau}\cdot\boldsymbol{\pi} + \bar{c}_1 \pi_3) S_\beta N\, F_{\mu\nu},\end{aligned} \tag{165}$$

with

$$\bar{d}_0, \bar{c}_0 = \mathcal{O}\left(e\delta_0 \frac{m_\pi^2}{M_{\not{T}}^2 M_{\mathrm{QCD}}}\right), \qquad \bar{d}_1, \bar{c}_1 = \mathcal{O}\left(e\delta_3 \frac{m_\pi^2}{M_{\not{T}}^2 M_{\mathrm{QCD}}}\right). \tag{166}$$

In contrast to the qCEDM, the isoscalar (isovector) qEDM generates isoscalar (isovector) nucleon-photon interactions, since the symmetry properties of the qEDM need not be mixed with chiral-symmetry breaking due to the quark charge.

In the case of the qEDM, long-range physics propagated by pions is suppressed by powers of α_{em}, and $\not{P}\not{T}$ observables are dominated by short-range nucleon-photon interactions. Since the operators in Eq. (165) contribute to the nucleon EDM only and not to the momentum-dependent part of the corresponding form factor, for the latter we need to construct electromagnetic operators with higher chiral index. It turns out that momentum dependence arises only at NNLO, so that we need to construct the Lagrangians with $\Delta_6 = 2, 3$.

At $\Delta_6 = 2$, the $\not{P}\not{T}$ electromagnetic operators that are not constrained by Lorentz invariance contain at least one pion,

$$\begin{aligned}
\mathcal{L}^{(2)}_{q,f=2\ \mathrm{em}} \quad = \quad & -\frac{i}{m_N}\bar{N}\left[\bar{d}_0\left(1-\frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right)+\bar{d}_1\left(\tau_3-\frac{2\pi_3}{F_\pi^2 D}\boldsymbol{\tau}\cdot\boldsymbol{\pi}\right)\right]S^\mu\mathcal{D}^\nu_{\perp\,-}N\,F_{\mu\nu} \\
& +\frac{i}{2F_\pi m_N D}\epsilon^{\mu\nu\alpha\beta}\bar{N}(\bar{c}_0\boldsymbol{\tau}\cdot\boldsymbol{\pi}+\bar{c}_1\pi_3)S_\beta\mathcal{D}_{\alpha\,-}N\,F_{\mu\nu} \\
& +\frac{1}{F_\pi}\bar{N}\left[\bar{\wp}_1\left(1-\frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right)(D^\mu\boldsymbol{\pi})\cdot\boldsymbol{\tau}+\bar{\rho}_1\left(D^\mu\pi_3-\frac{2\pi_3}{F_\pi^2 D}\boldsymbol{\pi}\cdot D^\mu\boldsymbol{\pi}\right)\right]N\,v^\nu F_{\mu\nu} \\
& +\frac{1}{F_\pi D}\bar{N}\left(\bar{\wp}_2\boldsymbol{\pi}\cdot\boldsymbol{\tau}+\bar{\rho}_2\pi_3\right)N\,v^\nu\partial^\mu F_{\mu\nu} \\
& +\frac{1}{F_\pi^2 D}\epsilon^{\alpha\beta\mu\nu}v_\alpha\bar{N}\left[\bar{\wp}_3(\boldsymbol{\pi}\cdot D_\beta\boldsymbol{\pi})+\bar{\rho}_3\pi_3\left(D_\beta\boldsymbol{\pi}\right)\cdot\boldsymbol{\tau}\right]N\,F_{\mu\nu} \\
& +\frac{1}{F_\pi^2 D}\bar{N}S^\mu\left[\bar{\wp}_4(\boldsymbol{\pi}\times D^\nu\boldsymbol{\pi})+\bar{\wp}_5(\boldsymbol{\pi}\times v\cdot D\boldsymbol{\pi})v^\nu\right]\cdot\boldsymbol{\tau}N\,F_{\mu\nu} \\
& +\frac{1}{F_\pi}\epsilon^{\alpha\beta\mu\nu}\,\bar{N}\left[(\bar{\rho}_4 D_\alpha\boldsymbol{\pi}+\bar{\rho}_5 v_\alpha v\cdot D\boldsymbol{\pi})\times\boldsymbol{\tau}\right]_i S_\beta N\left(\delta_{i3}-\frac{2\pi_3\pi_i}{F_\pi^2 D}\right)F_{\mu\nu}, \qquad (167)
\end{aligned}$$

where $\bar{\wp}_i$ ($\bar{\rho}_i$) originate from the isoscalar (isovector) qEDM, with

$$\bar{\wp}_i=\mathcal{O}\left(e\delta_0\frac{m_\pi^2}{M_{\not T}^2 M_{\mathrm{QCD}}^2}\right),\qquad \bar{\rho}_i=\mathcal{O}\left(e\delta_3\frac{m_\pi^2}{M_{\not T}^2 M_{\mathrm{QCD}}^2}\right). \qquad (168)$$

In Eq. (167) we already incorporated RPI, and the first two sets of interactions are recoil corrections to Eq. (165).

At $\Delta_6 = 3$, we encounter operators with two covariant derivatives or one insertion of the quark mass. Since it is unlikely that, at this order in the chiral series, operators containing pions are of any phenomenological use, we focus on terms without pions:

$$\begin{aligned}
\mathcal{L}^{(3)}_{q,f=2\,\mathrm{em}} \quad = \quad & -2\bar{N}\left(\delta\bar{d}_0+\delta\bar{d}_1\tau_3\right)S^\mu N\,v^\nu F_{\mu\nu} \\
& +\frac{1}{4m_N^2}\bar{N}\left(\bar{d}_0+\bar{d}_1\tau_3\right)S\cdot\mathcal{D}_{\perp-}\mathcal{D}^\mu_{\perp-}N\,v^\nu F_{\mu\nu} \\
& +\bar{N}\left(\bar{S}_0'+\bar{S}_1'\tau_3\right)\left(S\cdot\mathcal{D}_{\perp+}\mathcal{D}^\mu_{\perp+}+S^\mu\mathcal{D}^2_{\perp+}\right)N\,v^\nu F_{\mu\nu}+\ldots \qquad (169)
\end{aligned}$$

The first two terms are corrections to the isoscalar and isovector nucleon EDM due the quark mass, the third and fourth terms are relativistic corrections, and the operators with LECs $\bar{S}_0'$ and $\bar{S}_1'$ are the first qEDM contributions to, respectively, the isoscalar and isovector Schiff moments. The dots denote multi-pion components of the listed operators, and other operators that start at one pion, which we neglect. The LECs scale as

$$\begin{aligned}
\delta\bar{d}_0=\mathcal{O}\left(e\left(\delta_0+\varepsilon\delta_3\right)\frac{m_\pi^4}{M_{\not T}^2 M_{\mathrm{QCD}}^3}\right),\qquad & \delta\bar{d}_1=\mathcal{O}\left(e\left(\varepsilon\delta_0+\delta_3\right)\frac{m_\pi^4}{M_{\not T}^2 M_{\mathrm{QCD}}^3}\right), \\
\bar{S}_0'=\mathcal{O}\left(e\delta_0\frac{m_\pi^2}{M_{\not T}^2 M_{\mathrm{QCD}}^3}\right),\qquad & \bar{S}_1'=\mathcal{O}\left(e\delta_3\frac{m_\pi^2}{M_{\not T}^2 M_{\mathrm{QCD}}^3}\right). \qquad (170)
\end{aligned}$$

Here tadpoles are much smaller than the nucleon-photon interactions, and their removal does not affect the operators above.

8.4 χISs

The $\not{P}\not{T}$ chiral-invariant sources generate electromagnetic interactions that transform as $w\bar{I} \otimes e(I^\mu/6 + T^\mu_{34}/2)$ with chiral index $\Delta_6 = -1$,

$$\mathcal{L}^{(-1)}_{w,f=2\text{ em}} = -2\bar{N}\left\{\bar{d}_0 + \bar{d}_1\left[\tau_3 + \frac{2}{F_\pi^2 D}\left(\pi_3 \boldsymbol{\pi}\cdot\boldsymbol{\tau} - \boldsymbol{\pi}^2\tau_3\right)\right]\right\} S^\mu N\, v^\nu F_{\mu\nu}, \tag{171}$$

where the LECs scale as

$$\bar{d}^{(-1)}_{0,1} = \mathcal{O}\left(ew\frac{M_{\text{QCD}}}{M^2_{\not{T}}}\right). \tag{172}$$

Differently from qCEDM and FQLR, for chiral-invariant $\not{P}\not{T}$ sources the short-distance EDM operators have the same chiral index as the leading pion-nucleon coupling, Eq. (135). Since the latter contributes to the nucleon EDM only via loops, it follows that, for χISs, the nucleon EDM is mainly determined by short-distance physics.

A second consequence of the enhancement of short-distance vs. long-distance physics for χISs is that the nucleon EDFF does not depend on the momentum transfer at leading order in χPT. Momentum dependence only arises at NNLO, for which accuracy we need to consider the power-suppressed $\Delta_6 = 0, 1$ Lagrangian. We construct the complete chiral $\Delta_6 = 0$ Lagrangian,

$$\begin{aligned}
\mathcal{L}^{(0)}_{w,f=2\text{ em}} = & -\frac{i}{m_N}\bar{N}\left\{\bar{d}_0 + \bar{d}_1\left[\tau_3 + \frac{2}{F_\pi^2 D}\left(\pi_3\boldsymbol{\pi}\cdot\boldsymbol{\tau} - \boldsymbol{\pi}^2\tau_3\right)\right]\right\} S^\mu \mathcal{D}^\nu_{\perp -} N\, F_{\mu\nu} \\
& + \frac{\bar{\chi}_0}{F_\pi}(D^\mu\boldsymbol{\pi})\cdot\bar{N}\boldsymbol{\tau}N\, v^\nu F_{\mu\nu} \\
& + \frac{1}{F_\pi^2 D}\left[\bar{\chi}_1(\boldsymbol{\pi}\times D^\mu\boldsymbol{\pi})_3 + \bar{\chi}_2(\boldsymbol{\pi}\times v\cdot D\boldsymbol{\pi})_3 v^\mu\right]\bar{N}S^\nu N F_{\mu\nu} \\
& + \frac{\bar{\chi}_3}{F_\pi^2 D}\epsilon^{\alpha\beta\mu\nu}v_\alpha\bar{N}\left[(D_\beta\pi_3)\boldsymbol{\tau}\cdot\boldsymbol{\pi} - (\boldsymbol{\pi}\cdot D_\beta\boldsymbol{\pi})\tau_3\right]N\, F_{\mu\nu} \\
& + \frac{1}{F_\pi}\left\{\bar{\chi}_4(D^\mu\pi_i)\,\bar{N}Nv^\nu + \epsilon^{\alpha\beta\mu\nu}\bar{N}\left[\bar{\chi}_5(D_\alpha\boldsymbol{\pi}\times\boldsymbol{\tau})_i + \bar{\chi}_6(v\cdot D\boldsymbol{\pi}\times\boldsymbol{\tau})_i v_\alpha\right]S_\beta N\right\} \\
& \times\left[\delta_{i3} + \frac{2}{F_\pi^2 D}\left(\pi_3\pi_i - \boldsymbol{\pi}^2\delta_{i3}\right)\right]F_{\mu\nu} + \frac{\bar{\chi}_7}{F_\pi}\epsilon^{\alpha\beta\lambda\mu}v_\alpha\bar{N}(\boldsymbol{\tau}\times\boldsymbol{\pi})^3\mathcal{D}_{\beta +}S_\lambda N\, v^\nu F_{\mu\nu},
\end{aligned} \tag{173}$$

where the LECs scale as

$$\bar{\chi}_i = \mathcal{O}\left(\frac{ew}{M^2_{\not{T}}}\right). \tag{174}$$

At $\Delta_6 = 1$ we only construct the operators that start without pions, since operators with pions are of little phenomenological use. We have

$$\begin{aligned}
\mathcal{L}^{(1)}_{w,f=2\text{ em}} = & -2\bar{N}\left[\delta\bar{d}_0\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right) + \delta\bar{d}_1\left(\tau_3 - \frac{2\pi_3}{F_\pi^2 D}\boldsymbol{\pi}\cdot\boldsymbol{\tau}\right)\right] S^\mu N\, v^\nu F_{\mu\nu} \\
& -2\,\bar{\delta}d_1'\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right)\bar{N}\left[\tau_3 + \frac{2}{F_\pi^2 D}\left(\pi_3\boldsymbol{\pi}\cdot\boldsymbol{\tau} - \boldsymbol{\pi}^2\tau_3\right)\right] S^\mu N\, v^\nu F_{\mu\nu} \\
& + \frac{1}{4m_N^2}\bar{N}\left\{\bar{d}_0 + \bar{d}_1\left[\tau_3 + \frac{2}{F_\pi^2 D}\left(\pi_3\boldsymbol{\pi}\cdot\boldsymbol{\tau} - \boldsymbol{\pi}^2\tau_3\right)\right]\right\} S\cdot\mathcal{D}_{\perp -}\mathcal{D}^\mu_{\perp -}N\, v^\nu F_{\mu\nu} \\
& + \bar{N}\left(\bar{S}_0' + \bar{S}_1'\tau_3\right)\left(S\cdot\mathcal{D}_{\perp +}\mathcal{D}^\mu_{\perp +} + S^\mu\mathcal{D}^2_{\perp +}\right)N\, v^\nu F_{\mu\nu} + \dots
\end{aligned} \tag{175}$$

As for qEDM, the first operators (with LECs $\delta\bar{d}_{0,1}$ and $\bar{\delta}d'_1$) are corrections to the isoscalar and isovector nucleon EDMs proportional to the quark masses. The next interactions are relativistic corrections to Eq. (171). Finally, the last operators (with LECs $\bar{S}'_{0,1}$) are short-distance contributions to the first derivative of the nucleon EDFF, the Schiff moment. The scaling of the LECs is

$$\delta\bar{d}_{0,1} = \mathcal{O}\left(e(1+\varepsilon)w\frac{m_\pi^2}{M_{\not{T}}^2 M_{\mathrm{QCD}}}\right), \qquad \delta\bar{d}'_1 = \mathcal{O}\left(ew\frac{m_\pi^2}{M_{\not{T}}^2 M_{\mathrm{QCD}}}\right), \qquad \bar{S}'_{0,1} = \mathcal{O}\left(\frac{ew}{M_{\not{T}}^2 M_{\mathrm{QCD}}}\right). \tag{176}$$

The leading nucleon-photon interactions in Eq. (171) are chiral invariant or transform as the 34 component of an antisymmetric tensor. In both cases, they are not affected by the field redefinition in Eq. (91). Only the NNLO operators $\delta\bar{d}_{0,1}$ and $\delta\bar{d}'_1$, which are proportional to the quark masses, are affected by the elimination of the tadpoles, in a way that leads to the shift $w \to w(1+\varepsilon^2)$.

9 Nucleon EDM from the FQLR

The electromagnetic interactions of the previous section allow the calculation of EDFFs. In fact, the chiral Lagrangians from the qEDM, qCEDM, and χISs derived in this article have already been used to calculate the $\not{P}\not{T}$ form factors of the nucleon and several light nuclei in Refs. [43, 33, 34, 36, 37]. However, in these references the contributions from the FQLR operator were omitted. In this section we update our previous work by giving the FQLR contributions to the nucleon EDM to NLO. Nuclear issues are discussed in the next section.

The nucleon EDFF, following Refs. [29, 43], is decomposed as

$$F_i(Q^2) = d_i - S'_i Q^2 + H_i(Q^2), \tag{177}$$

where d_i is the isospin-i component of the EDM, S'_i is the corresponding Schiff moment [28], and $H_i(Q^2)$ accounts for the remaining dependence on $Q^2 = -q^2 > 0$, q being the outgoing momentum of the photon. The EDFF of the proton (neutron) is $F_0 + F_1$ ($F_0 - F_1$).

The LO $\not{P}\not{T}$ pion-nucleon and nucleon-photon interactions from the FQLR are very similar to those from an isovector qCEDM, apart from interactions involving multiple pions, which contribute at higher orders. Therefore, the LO nucleon EDFF from the FQLR is of identical form as that from the isovector qCEDM calculated in Ref. [43], but, of course, with different scalings for the LECs. The LO nucleon EDFF gets contributions from the short-range interactions in Eq. (163) and from one-loop diagrams involving the LO isoscalar $\not{P}\not{T}$ pion-nucleon vertex $\bar{g}_0$ in Eq. (120) (*cf.* Ref. [4, 28, 29]). At NLO, there are additional one-loop contributions, involving both $\bar{g}_0$ and the isovector $\not{P}\not{T}$ vertex $\bar{g}_1$, and subleading couplings in the PT-even Lagrangian (*cf.* Refs. [31, 40, 33]). One has in principle to consider also the effects of the three-pion coupling $\bar{\Delta}_{\mathrm{LR}}^{(-4)}$ in Eq. (100), which contributes to the nucleon EDM via two-loop diagrams. However, it is easy to see that all the two-loop diagrams vanish because of their isospin structure, so that the nucleon EDM at NLO stemming from the FQLR is identical to that generated by the qCEDM. In the evaluation of loop diagrams we use dimensional regularization, as in Sec. 7.

The leading loop diagrams do not generate an isoscalar EDFF, which is therefore purely tree-level and static at LO. The first non-analytic contribution to the isoscalar EDM arises at NLO, and it has both $\bar{g}_0$ and $\bar{g}_1$ pieces. At the same order one also finds the first momentum dependence of the isoscalar EDFF, which is proportional to the hadronic part of the nucleon mass splitting, δm_N in Eq. (67), and the isoscalar coupling $\bar{g}_0$. The isoscalar EDFF to NLO is given by

$$\begin{aligned} d_0 &= \bar{d}_0 + \pi \frac{eg_A\bar{g}_0}{(2\pi F_\pi)^2}\left[\frac{3m_\pi}{4m_N}\left(1+\frac{\bar{g}_1}{3\bar{g}_0}\right) - \frac{\delta m_N}{m_\pi}\right], && (178)\\ S_0' &= -\frac{\pi}{12m_\pi^2}\frac{eg_A\bar{g}_0}{(2\pi F_\pi)^2}\frac{\delta m_N}{m_\pi}, && (179)\\ H_0(Q^2) &= -\frac{\pi}{5}\frac{eg_A\bar{g}_0}{(2\pi F_\pi)^2}\frac{\delta m_N}{m_\pi}\,h_0^{(1)}\left(\frac{Q^2}{4m_\pi^2}\right), && (180) \end{aligned}$$

with the function

$$h_0^{(1)}(x) \equiv 5\left(\frac{1}{\sqrt{x}}\arctan\sqrt{x} - 1 + \frac{x}{3}\right), \qquad (181)$$

such that $h_0^{(1)}(x) = x^2 + \mathcal{O}(x^3)$ for $x \ll 1$.

In contrast, for the isovector EDM loop diagrams do renormalize short-distance operators, in addition to generating a non-trivial momentum dependence. Instead of the nucleon mass splitting from the quark masses, there is a contribution from the electromagnetic pion mass splitting $\breve{\delta} m_\pi^2$ in Eq. (69). To NLO, the isovector EDM is found to be

$$d_1 = \bar{d}_1 + \bar{d}_1' + \frac{eg_A\bar{g}_0}{(2\pi F_\pi)^2}\left[L - \ln\frac{m_\pi^2}{\mu^2} + \frac{5\pi}{4}\frac{m_\pi}{m_N}\left(1+\frac{\bar{g}_1}{5\bar{g}_0}\right) - \frac{\breve{\delta} m_\pi^2}{m_\pi^2}\right], \qquad (182)$$

while the momentum dependence is given by

$$\begin{aligned} S_1' &= \frac{1}{6m_\pi^2}\frac{eg_A\bar{g}_0}{(2\pi F_\pi)^2}\left[1 - \frac{5\pi}{4}\frac{m_\pi}{m_N} - \frac{\breve{\delta} m_\pi^2}{m_\pi^2}\right], && (183)\\ H_1(Q^2) &= \frac{4}{15}\frac{eg_A\bar{g}_0}{(2\pi F_\pi)^2}\left[h_1^{(0)}\left(\frac{Q^2}{4m_\pi^2}\right) - \frac{7\pi}{8}\frac{m_\pi}{m_N}\,h_1^{(1)}\left(\frac{Q^2}{4m_\pi^2}\right) - \frac{2\breve{\delta} m_\pi^2}{m_\pi^2}\,\breve{h}_1^{(1)}\left(\frac{Q^2}{4m_\pi^2}\right)\right]. && (184) \end{aligned}$$

Here the functions

$$\begin{aligned} h_1^{(0)}(x) &\equiv -\frac{15}{4}\left[\sqrt{1+\frac{1}{x}}\ln\left(\frac{\sqrt{1+\frac{1}{x}}+1}{\sqrt{1+\frac{1}{x}}-1}\right) - 2\left(1+\frac{x}{3}\right)\right], && (185)\\ h_1^{(1)}(x) &\equiv -\frac{1}{7}\left[3(1+2x)\,h_1^{(0)}(x) - 10x^2\right], && (186)\\ \breve{h}_1^{(1)}(x) &\equiv -\frac{1}{4(1+x)}\left(h_1^{(0)}(x) - 5x^2\right) && (187) \end{aligned}$$

are defined so that they satisfy $h_1^{(i)}(x \ll 1) = x^2 + \mathcal{O}(x^3)$.

The dependence on the arbitrary scale μ from the loops is compensated by the counterterm $\bar{d}_1 + \bar{d}_1'$. In fact, the loop contributions cannot be separated from the short-range pieces in a model-independent way. However, one does not expect cancellations between the non-analytic dependence in the pion mass and the analytic dependence of the short-range part. Taking the long-range part in Eq. (182) at $\mu = m_N$ as an estimate for the neutron EDM, and using relation (122) with $\delta m_N \simeq 2.3$ MeV [60],

$$|d_n| \gtrsim \frac{e g_A \delta m_N |\bar{\Delta}_{\mathrm{LR}}^{(-4)}|}{2\pi^2 m_\pi^2 F_\pi^2} \ln \frac{m_N}{m_\pi} \simeq 0.003 \frac{|\bar{\Delta}_{\mathrm{LR}}^{(-4)}|}{F_\pi^2}\, e\,\mathrm{fm}. \tag{188}$$

The expected scaling of $\bar{\Delta}_{\mathrm{LR}}^{(-4)}$ in Eq. (79) with $M_{\mathrm{QCD}} \sim 2\pi F_\pi$ gives $|d_n| \sim 0.8\,\mathrm{GeV}\, e|\xi|/M_{\not T}^2$, which is about two thirds of the direct NDA estimate for $\bar{d}_1$ and $\bar{d}_1'$, Eq. (164). We therefore expect the latter to provide a better estimate for the neutron EDM, and

$$|d_n| \sim 1.2\,\mathrm{GeV}\,\frac{e|\xi|}{M_{\not T}^2}. \tag{189}$$

Note that the neutron EDM was computed from general $\not P \not T$ four-quark operators in Ref. [48], where it was found to be $|d_n| \simeq 0.15\,\mathrm{GeV}\, e\,|C_4|$ in terms of C_4, the coupling constant of the four-quark operators. We compare to our result by writing $C_4 = (4\pi)^2 \xi / M_{\not T}^2$, and find that it is about an order of magnitude larger than our estimate. This discrepancy can be traced [63] to the use of a relativistic formalism in Ref. [48]. In this reference, a large contribution to the nucleon EDM comes from pion loops in which the photon couples to the nucleon via the nucleon anomalous magnetic moment, while $\not P \not T$ is provided by the coupling $\bar{g}_1$. While a chiral logarithm $\log m_\pi^2/m_N^2$ appears with a suppression of m_π^2/m_N^2, the contributions from loop momenta $|\vec{k}| \sim m_N$ give rise to a large EDM, which does not have such a suppression. In the heavy-baryon formalism, the nucleon magnetic moment does contribute to the nucleon EDM, but only at NNLO, and should reproduce the $\log m_\pi^2$ dependence of the relativistic calculation [63]. Unfortunately, χPT cannot be trusted for momenta where the nucleon is relativistic. Thus, we interpret the result of Ref. [48] as a model-dependent estimate of the size of the counterterms $\bar{d}_1$, $\bar{d}_1'$ and $\bar{d}_0$, which, in this calculation, appears to be somewhat larger than NDA.

10 Nuclear effects

So far we have focused on interactions involving at most one nucleon. Power counting is more complicated for processes involving more than one nucleon: infrared enhancements due to purely nucleonic states require a resummation of leading interactions [23], and this resummation in turn leads to the enhancement of certain *CP*-even operators that one would naively think are of high order [64]. Thus, the chiral index Δ in Eq. (62) is not particularly useful due to non-perturbative renormalization.

On the other hand, $\not P \not T$ operators should be treated perturbatively and are likely unaffected by this subtlety. It is convenient then to continue to organize them according to the index Δ_6 in Eq. (73), where f can now take even values larger than 2. Since the index increases with increasing f, the operators most likely to be relevant have $f = 4$. The most important four-nucleon $\not P \not T$ operators have been constructed in Ref. [35]. We give here some of the details for dimension-six sources. The consequences for the $\not P \not T$ nuclear potential induced by the qCEDM, χISs and qEDM can be found in Ref. [35]. Ref. [35] did not discuss the $\not P \not T$ potential stemming from the FQLR operator. In this section, after constructing the short-range NN and $NN\pi$ operators induced by each $\not P \not T$ source, we remedy this omission.

10.1 χISs

While in the pion-nucleon sector one needs at least either two derivatives or one insertion of the quark mass to generate operators from the χISs, in the nucleon-nucleon sector one derivative is enough, and it need not include a pion field:

$$\mathcal{L}^{(-1)}_{w,f=4} = \bar{C}_1 \bar{N}N\, \partial_\mu(\bar{N}S^\mu N) + \bar{C}_2 \bar{N}\boldsymbol{\tau}N \cdot \mathcal{D}_\mu(\bar{N}S^\mu \boldsymbol{\tau} N), \tag{190}$$

with scalings

$$\bar{C}_{1,2} = \mathcal{O}\left(w \frac{M_{\mathrm{QCD}}}{F_\pi^2 M_{\not{T}}^2}\right). \tag{191}$$

We see that for the χISs, the chiral index of the LO $f = 4$ LECs is the same as that of the LO $f = 2$ LECs, in contrast with sources that are chiral breaking. Thus, whereas for $\bar{\theta}$, qCEDM, and FQLR the potential is dominated by pion exchange (and photon exchange for qEDM), the potential for the χISs gets contributions of the same order from pion exchange and short-range NN interactions [35].

The first terms linear in the pion field appear one order higher,

$$\mathcal{L}^{(0)}_{w,f=4} = \frac{\mathcal{D}_\mu \boldsymbol{\pi}}{F_\pi} \cdot \left(\bar{G}_1 \bar{N}S^\mu \boldsymbol{\tau}N\, \mathcal{D}_\nu(\bar{N}S^\nu N) + \bar{G}_2 \bar{N}S^\mu N\, \mathcal{D}_\nu(\bar{N}\boldsymbol{\tau}S^\nu N)\right), \tag{192}$$

with scalings

$$\bar{G}_{1,2} = \mathcal{O}\left(\frac{w}{F_\pi^2 M_{\not{T}}^2}\right). \tag{193}$$

When the pion is attached to another nucleon, a three-nucleon force results.

The interactions above are not affected by tadpole removal because the operators are chiral invariant. At NNLO we find additional $\not{P}\not{T}$ nuclear interactions, which we do not construct. At this order the first contributions to the isovector $\not{P}\not{T}$ nucleon-nucleon interactions appear as well, originating from an insertion of the quark mass difference.

10.2 qCEDM

Because it is chiral breaking, the qCEDM generates pion interactions at lower order than short-range, purely nucleonic interactions. At $\Delta_6 = 0$,

$$\begin{aligned} \mathcal{L}^{(0)}_{\tilde{q},f=4} &= -\frac{1}{F_\pi D}\boldsymbol{\pi} \cdot \left(\bar{\gamma}_1 \bar{N}\boldsymbol{\tau}N\, \bar{N}N + \bar{\gamma}_2 \bar{N}\boldsymbol{\tau}S_\mu N\, \bar{N}S^\mu N\right) \\ &\quad -\frac{\pi_3}{F_\pi D}\left(\bar{\gamma}_3 \bar{N}N\, \bar{N}N + \bar{\gamma}_4 \bar{N}S_\mu N\, \bar{N}S^\mu N\right), \end{aligned} \tag{194}$$

where, after tadpole extermination, the LECs scale as

$$\bar{\gamma}_{1,2} = \mathcal{O}\left(\left(\tilde{\delta}_0 + \varepsilon\tilde{\delta}_3\right)\frac{m_\pi^2}{F_\pi^2 M_{\not{T}}^2}\right), \qquad \bar{\gamma}_{3,4} = \mathcal{O}\left(\tilde{\delta}_3 \frac{m_\pi^2}{F_\pi^2 M_{\not{T}}^2}\right). \tag{195}$$

One order down, we find the first short-range interactions contributing directly to nucleon-nucleon scattering,

$$\begin{aligned} \mathcal{L}^{(1)}_{\tilde{q},f=4} &= \left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right)\left[\bar{C}_1 \bar{N}N\, \partial_\mu(\bar{N}S^\mu N) + \bar{C}_2 \bar{N}\boldsymbol{\tau}N \cdot \mathcal{D}_\mu(\bar{N}S^\mu \boldsymbol{\tau}N)\right] \\ &\quad + \left(\delta_{3i} - \frac{2\pi_3\pi_i}{F_\pi^2 D}\right)\left[\bar{C}_3 \bar{N}\tau_i N\, \partial_\mu\left(\bar{N}S^\mu N\right) + \bar{C}_4 \bar{N}N\, \mathcal{D}_\mu\left(\bar{N}\tau_i S^\mu N\right)\right], \end{aligned} \tag{196}$$

with the scaling

$$\bar{C}_{1,2} = \mathcal{O}\left(\left(\tilde{\delta}_0 + \varepsilon\tilde{\delta}_3\right)\frac{m_\pi^2}{F_\pi^2 M_{\not T}^2 M_{\mathrm{QCD}}}\right), \qquad \bar{C}_{3,4} = \mathcal{O}\left(\tilde{\delta}_3\frac{m_\pi^2}{F_\pi^2 M_{\not T}^2 M_{\mathrm{QCD}}}\right). \tag{197}$$

Here, as before, the LECs of the operators induced by the isoscalar qCEDM get a contribution from the isovector qCEDM as well, proportional to ε. At this order, for all sources, there appear operators that start with one or more pions. We do not list them.

A comparison between the pion-nucleon and nuclear sectors shows that for qCEDM the most important $\not P \not T$ pion-nucleon interactions are larger by a factor M_{QCD}^2/Q^2 than the short-range $\not P \not T$ nucleon-nucleon interactions, implying that for these sources the $\not P \not T$ nuclear potential is dominated by pion exchange [35]. This observation justifies, *a posteriori*, the assumption often made in the literature, that $\not P \not T$ nuclear observables can be calculated solely from $\not P \not T$ pion exchange. However, as we have seen, for χISs this assumption is not valid.

10.3 FQLR

As we saw already in the pion and pion-nucleon sectors, the isovector qCEDM and FQLR generate very similar interactions, if one neglects the more complicated pion structure of the FQLR.

Here again,

$$\mathcal{L}^{(-2)}_{\mathrm{LR},f=4} = \mathcal{L}^{(0)}_{\tilde{q},f=4} + \ldots, \tag{198}$$

where "…" represent interactions with more pion fields, which are of little importance. The LECs scale as

$$\bar{\gamma}_{1,2} = \mathcal{O}\left(\varepsilon\xi\frac{M_{\mathrm{QCD}}^2}{F_\pi^2 M_{\not T}^2}\right), \qquad \bar{\gamma}_{3,4} = \mathcal{O}\left(\xi\frac{M_{\mathrm{QCD}}^2}{F_\pi^2 M_{\not T}^2}\right) \tag{199}$$

after tadpole removal. Similarly,

$$\mathcal{L}^{(-1)}_{\mathrm{LR},f=4} = \mathcal{L}^{(1)}_{\tilde{q},f=4} + \ldots, \tag{200}$$

with

$$\bar{C}_{1,2} = \mathcal{O}\left(\varepsilon\xi\frac{M_{\mathrm{QCD}}}{F_\pi^2 M_{\not T}^2}\right), \qquad \bar{C}_{3,4} = \mathcal{O}\left(\xi\frac{M_{\mathrm{QCD}}}{F_\pi^2 M_{\not T}^2}\right). \tag{201}$$

As for qCEDM, one-pion exchange is more important than these short-range interactions [35], which we do not pursue further.

10.4 qEDM

The qEDM, while sharing with qCEDM and FQLR the property that interactions involving at least one pion dominate [35], produces a richer isospin structure for the short-range nucleon-nucleon interactions, due to the integration of hard photon.

At $\Delta_6 = 3$ we find interactions that transform as the tensor product $(-d_0 V_4 + d_3 W_3) \otimes e(I^\mu/6 + T^\mu_{34}/2)$,

$$\begin{aligned}
\mathcal{L}^{(3)}_{q,f=4} &= -\frac{1}{F_\pi D}\boldsymbol{\pi}\cdot\left(\bar\gamma_1 \bar N\boldsymbol{\tau}N\,\bar NN + \bar\gamma_2 \bar N\boldsymbol{\tau}S_\mu N\,\bar N S^\mu N\right) \\
&\quad -\frac{\pi_3}{F_\pi D}\left(\bar\gamma_3 \bar NN\,\bar NN + \bar\gamma_4 \bar N S_\mu N\,\bar N S^\mu N\right) \\
&\quad -\frac{\pi_3}{F_\pi D}\left[\delta_{3i} + \frac{2}{F_\pi^2 D}\left(\pi_3\pi_i - \boldsymbol{\pi}^2\delta_{3i}\right)\right]\left[\bar\gamma_5 \bar N\tau_i N\bar NN + \bar\gamma_6 \bar N\tau_i S_\mu N\bar N S^\mu N\right],
\end{aligned} \tag{202}$$

with

$$\bar\gamma_{1,2,3,4} = \mathcal{O}\left((\delta_0+\delta_3)\,\frac{\alpha_{\rm em}}{4\pi}\frac{m_\pi^2}{F_\pi^2 M_{\not T}^2}\right), \qquad \bar\gamma_{5,6} = \mathcal{O}\left(\delta_3\frac{\alpha_{\rm em}}{4\pi}\frac{m_\pi^2}{F_\pi^2 M_{\not T}^2}\right). \tag{203}$$

One order higher we find

$$\begin{aligned}
\mathcal{L}^{(4)}_{q,f=4} &= \left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right)\left[\bar C_1 \bar NN\,\partial_\mu(\bar N S^\mu N) + \bar C_2 \bar N\boldsymbol{\tau}N\cdot\mathcal{D}_\mu(\bar N S^\mu\boldsymbol{\tau}N)\right] \\
&\quad + \left(\delta_{3i} - \frac{2\pi_3\pi_i}{F_\pi^2 D}\right)\left[\bar C_3 \bar N\tau_i N\,\partial_\mu\left(\bar N S^\mu N\right) + \bar C_4 \bar NN\,\mathcal{D}_\mu\left(\bar N\tau_i S^\mu N\right)\right] \\
&\quad + \left\{\left(1 - \frac{2\boldsymbol{\pi}^2}{F_\pi^2 D}\right)\left[\bar C_5 \bar N\tau_l N\,\partial_\mu\left(\bar N S^\mu N\right) + \bar C_6 \bar NN\,\mathcal{D}_\mu\left(\bar N\tau_l S^\mu N\right)\right]\right. \\
&\quad \left. + \bar C_7\left(\delta_{3k} - \frac{2\pi_3\pi_k}{F_\pi^2 D}\right)\bar N\tau_k N\,\mathcal{D}_\mu\left(\bar N S^\mu \tau_l N\right)\right\}\left[\delta_{l3} + \frac{2}{F_\pi^2 D}\left(\pi_3\pi_l - \boldsymbol{\pi}^2\delta_{3l}\right)\right] \\
&\quad + \ldots,
\end{aligned} \tag{204}$$

where we have ignored operators that start with one or more pions, and

$$\begin{aligned}
&\bar C_{1,2,3,4} = \mathcal{O}\left((\delta_0+\delta_3)\frac{\alpha_{\rm em}}{4\pi}\frac{m_\pi^2}{F_\pi^2 M_{\not T}^2 M_{\rm QCD}}\right), \qquad \bar C_{5,6} = \mathcal{O}\left(\delta_0\frac{\alpha_{\rm em}}{4\pi}\frac{m_\pi^2}{F_\pi^2 M_{\not T}^2 M_{\rm QCD}}\right), \\
&\bar C_7 = \mathcal{O}\left(\delta_3\frac{\alpha_{\rm em}}{4\pi}\frac{m_\pi^2}{F_\pi^2 M_{\not T}^2 M_{\rm QCD}}\right).
\end{aligned} \tag{205}$$

Since these operators are all suppressed by $\alpha_{\rm em}/4\pi$, their phenomenological impact is minimal, and we do not construct operators with higher chiral index.

10.5 The $\not P\not T$ potential from the FQLR

In Ref. [35] it was found that for the qCEDM the two-body $\not P\not T$ potential is dominated by one-pion exchange (OPE), where one of the pion couplings is a $\not P\not T$ coupling from Sec. 6. Short-range nucleon-nucleon interactions play a role at NNLO. On the other hand, for χISs short-range interactions appear at LO, while for the qEDM both pion and photon exchange contribute to the LO potential, which, however, does not usually play an important role in the calculation of EDMs. For qCEDM, qEDM and χISs, few-body $\not P\not T$ forces are suppressed. For FQLR, we will see that the forces are somewhat different. We will show that, in LO, the two-nucleon $\not P\not T$ potential is dominated by OPE and it is very similar to the potential induced by the qCEDM. However, the first loop correction appears already at NLO. The most striking feature of the



Figure 3: Contribution from the purely mesonic $\not{P}\not{T}$ interaction to the three-nucleon potential. Notation as in Fig. 2.

FQLR is, however, that the three-pion vertex $\bar{\Delta}^{(-4)}_{\rm LR}$ induces a three-nucleon $\not{P}\not{T}$ force at LO, which impacts the calculation of the EDMs of ^{3}He and ^{3}H.

We write the potential in terms of the spin, isospin, and incoming (outgoing) momentum of nucleon i, $\vec{\sigma}^{(i)}$, $\boldsymbol{\tau}^{(i)}$, and $\vec{p}_i$ ($\vec{p}^{\,\prime}_i$) respectively. We denote the transferred momenta by $\vec{q}_i = \vec{p}_i - \vec{p}^{\,\prime}_i$. In the two-nucleon case, $\vec{q}_2 = -\vec{q}_1$, while for three nucleons, $\vec{q}_3 = -(\vec{q}_1 + \vec{q}_2)$. The coordinate-space version of the potential can be obtained straightforwardly following the procedure of, for example, Ref. [35].

At LO in χPT, the $\not{P}\not{T}$ two-nucleon potential is given by OPE, with T violation provided by the pion-nucleon couplings of Sec. 6.3. The LO two-nucleon potential is thus identical to that generated by the qCEDM, and it is given by

$$\begin{aligned} V^{(-3)}_{2,\rm FQLR}(\vec{q}_1) &= i\frac{g_A}{F_\pi^2}\frac{\vec{q}_1}{\vec{q}_1^{\,2}+m_\pi^2}\cdot\Big\{\bar{g}_0\,\boldsymbol{\tau}^{(1)}\cdot\boldsymbol{\tau}^{(2)}\left(\vec{\sigma}^{(1)}-\vec{\sigma}^{(2)}\right) \\ &\quad +\frac{(\bar{g}_1+\bar{g}_1')}{2}\left[\left(\tau_3^{(1)}+\tau_3^{(2)}\right)\left(\vec{\sigma}^{(1)}-\vec{\sigma}^{(2)}\right)+\left(\tau_3^{(1)}-\tau_3^{(2)}\right)\left(\vec{\sigma}^{(1)}+\vec{\sigma}^{(2)}\right)\right]\Big\} \end{aligned} \quad (206)$$

Since the couplings $\bar{g}_0$ and $\bar{g}_1'$ are not independent, this potential depends on two low-energy constants, which we can choose to be $\bar{g}_1$ and $\bar{g}_1'$ using Eq. (123). The isoscalar piece of the potential is suppressed by the smallness of the ratio of the nucleon mass difference and the sigma term. Unless $\bar{g}_1$ is unnaturally tuned to cancel the contribution of $\bar{g}_1'$, the two-nucleon potential is mainly isovector.

In Sec. 7.4 we showed that the isovector PNFF F_2 receives loop corrections, and in particular non-analytic momentum dependence, at NLO. These corrections are proportional to the three-pion vertex $\bar{\Delta}^{(-4)}_{\rm LR}$, which is related to $\bar{g}_1'$ by Eq. (122), and are enhanced by a factor of π with respect to NDA. Their contribution to the isovector potential can thus be sizable, and, furthermore, it comes with a different momentum dependence with respect to the LO OPE.

The deuteron EDM and MQM induced by the FQLR at LO can therefore be read off the calculations for the qCEDM of Refs. [36, 37], after one accounts for the suppression of $\bar{g}_0$. At NLO, the deuteron EDM and MQM receive contributions from the three-pion vertex $\bar{\Delta}^{(-4)}_{\rm LR}$, and the NLO corrections could be important. While we postpone a detailed analysis of the $\bar{\Delta}^{(-4)}_{\rm LR}$ correction to the deuteron EDM, we expect the main qualitative conclusion of Ref. [36] to continue to hold —that is, that the deuteron EDM induced by an isospin-breaking $\not{P}\not{T}$ source is expected to be significantly larger than the isoscalar nucleon EDM.

The three-pion vertex $\bar{\Delta}^{(-4)}_{\rm LR}$ has even more striking consequences in the three-nucleon system. This interaction induces a three-nucleon potential, depicted in Fig. 3, which in the power counting of Ref. [36] enters at the same order as the LO two-nucleon potential. The three-nucleon potential is of the form

$$V_{3,\mathrm{FQLR}}^{(-3)}(\vec{q}_1,\vec{q}_2) = \frac{2g_A^3\bar{\Delta}_{\mathrm{LR}}^{(-4)}}{F_\pi^4}\left(\tau_3^{(1)}\,\boldsymbol{\tau}^{(2)}\cdot\boldsymbol{\tau}^{(3)}+\tau_3^{(2)}\,\boldsymbol{\tau}^{(1)}\cdot\boldsymbol{\tau}^{(3)}+\tau_3^{(3)}\,\boldsymbol{\tau}^{(1)}\cdot\boldsymbol{\tau}^{(2)}\right) \times\frac{\vec{\sigma}^{(1)}\cdot\vec{q}_1\;\vec{\sigma}^{(2)}\cdot\vec{q}_2\;\vec{\sigma}^{(3)}\cdot(\vec{q}_1+\vec{q}_2)}{(\vec{q}_1^{\,2}+m_\pi^2)(\vec{q}_2^{\,2}+m_\pi^2)[(\vec{q}_1+\vec{q}_2)^2+m_\pi^2]}. \tag{207}$$

The size of the contributions from Eq. (207) can be compared to those from the LO two-nucleon potential (206) by using the power-counting rules outlined in Ref. [36]. These rules indicate that the three-body contributions scale exactly as the LO two-body contributions. Equation (207) provides a correction to the helion and triton EDMs, which, by power counting, is of the same size as the two-body terms calculated in Ref. [36]. While we expect that the main qualitative conclusions of Ref. [36] —that for an isospin-breaking source the EDMs of helion and triton are significantly different from the EDMs of their constituents— to hold, we leave a quantitative analysis to future work.

Notice that, as always, power counting provides only a guide for what should be included in a calculation. In particular, the power counting for few-nucleon forces has not been significantly probed. The one adopted in Ref. [36], which is in line with the current understanding of the two-body system, agrees with the one in Ref. [65]. If, instead, the power counting of Refs. [23, 66] is employed, then Eq. (207) becomes instead an NLO correction.

The discussion of the two- and three-nucleon potentials shows that for $\not{P}\not{T}$ observables in light nuclei the three-pion vertex $\bar{\Delta}_{\mathrm{LR}}^{(-4)}$, which survives tadpole extermination because of the tensor nature of the FQLR, plays an important role. Another handle to disentangle the qCEDM and the FQLR could be provided by the measurement of the T-odd correlation coefficient in β decay, D [22]. Indeed, the Ξ_1 operator in Eq. (24), which generates at tree level the FQLR operator, also contributes to an operator that couples right-handed quarks to left-handed electron and neutrino, with coefficient of $\mathcal{O}(\Xi_1)$. The qCEDM contributions to the same operator are suppressed by G_F due to the need for the exchange of a W boson and, because of an extra chirality flip, by an insertion of the light-quark mass. Therefore, the contribution of the qCEDM to the operator scales as $\tilde{d}_{0,3}G_F\bar{m}/4\pi \sim G_F\bar{m}^2\tilde{\delta}_{0,3}/M_{\not{T}}^2$, which is suppressed by a factor $\bar{m}^2/M_W^2$. The size of the D coefficient with respect to nuclear EDMs is therefore very different for isovector qCEDM and FQLR operators. In case that the observation of nucleon and deuteron EDM point to an isospin-breaking $\not{P}\not{T}$ source, the measurement of D could help to discriminate between these two operators.

11 Discussion and conclusion

As we have seen, the chiral structure of the various $\not{P}\not{T}$ sources is influential in the form and expected magnitude of the low-energy $\not{P}\not{T}$ interactions. Even without going into detailed results for $\not{P}\not{T}$ hadronic observables we can draw some qualitative conclusions by looking at the Lagrangian that we constructed. In Ref. [32] it was found that for the Standard Model $\bar{\theta}$ term, all LECs are proportional to negative powers of the scale M_{QCD}. The reasons for this are twofold: the $\bar{\theta}$ term (i) can be seen as a complex quark mass term, which brings in at least one power of m_π^2, and (ii) transforms, once vacuum stability is imposed, as the fourth component of an $SO(4)$ vector, which means that a pion tadpole can be eliminated. Just like isospin violation

[54], time reversal is therefore an accidental symmetry in the SM, in the sense that it would be somewhat suppressed (by at least one power of $m_\pi/M_{\rm QCD}$) even if $\bar\theta$ were not small.

Extending the analysis to the dimension-six operators, we see that positive powers of $M_{\rm QCD}$ do appear, but they are of course overcompensated by two negative powers of the much larger $M_{\not T}$. The main result of tadpole extermination is that the vacuum misalignment signaled by the pion tadpoles causes the isoscalar operators, like $\bar g_0$ or $\bar C_{1,2}$, to receive additional contributions from isospin-breaking sources, like the isovector qCEDM and the FQLR operator, of the same importance as the contributions of isoscalar sources. These contributions can be schematically obtained by replacing $\tilde\delta_0$ before tadpole removal with $\tilde\delta_0 + \varepsilon\tilde\delta_3 + \varepsilon\xi M^2_{QCD}/m^2_\pi$ in the power counting estimates of isoscalar operators. Shifts in isovector quantities are inconsequential in the sense that they do not change the dependence of the LECs on ξ, $\tilde\delta_{0,3}$, and w. It is in the case of FQLR that tadpole rotation is most important because, first, FQLR does not have an isoscalar piece and, second, a relatively important three-pion vertex survives.

We notice that the different chiral properties and field content of the $\bar\theta$ term, qCEDM, and FQLR on the one side, and qEDM and χISs on the other, imply very different relations between long-distance and short-distance $\not P\not T$ effects. The $\bar\theta$ term, qCEDM, and FQLR all violate chiral symmetry and thus generate $\not P\not T$ pion-nucleon interactions in which the pion couples to the nucleon non-derivatively. As a consequence, the first $\not P\not T$ pion-nucleon couplings appear in the Lagrangian two orders before short-range contributions to the nucleon EDM. For the nucleon EDM and EDFF, this fact implies that even though pion-nucleon $\not P\not T$ couplings can only contribute to the nucleon EDM via loops, which bring in a $m^2_\pi/(2\pi F_\pi)^2$ suppression, they are still as important as the short-distance operators [29, 43]. For light nuclei, such as the deuteron [62, 34] and helion [67, 36], the most important contribution to $\not P\not T$ electromagnetic moments comes from the $\not P\not T$ OPE potential —which causes the nucleus wavefunction to mix with states of different parity— unless the admixed component has quantum numbers that cause the dipole matrix element to vanish. The application of chiral EFT to study effects of $\bar\theta$ term and qCEDM in systems with $A \geq 2$ nucleons is thus particularly promising, since $\not P\not T$ observables are likely to depend on few LECs from the $f = 2$ $\not P\not T$ Lagrangian. Once these constants are fixed in experiments, one is in the position to make testable, model-independent predictions. For the FQLR, there appear additional complications for EDMs of nuclei due to a $\not P\not T$ three-pion interaction, which induces an NLO correction to the two-nucleon potential and, more importantly, a LO three-nucleon potential. The contributions of this three-nucleon force to light-nuclear EDMs have, so far, not been calculated.

For the chiral-invariant $\not P\not T$ sources, instead, the pion-nucleon $\not P\not T$ coupling appear in the Lagrangian at the same order as short-distance nucleon EDM operators. This happens because it is not possible to write a $\not P\not T$ chiral-invariant pion-nucleon coupling with only one derivative. The first chiral-invariant $\not P\not T$ pion-nucleon coupling must have two derivatives, while nonderivative couplings can be generated by considering the combined effects of chiral-invariant $\not P\not T$ sources and the chiral-breaking quark mass; in any case, pion-nucleon couplings receive a further suppression of $Q^2/M^2_{\rm QCD}$. It should be noted that this difficulty does not affect the nucleon-photon and nucleon-nucleon sectors, where operators with a minimal number of derivatives can be constructed. The consequence for the nucleon EDM is that it is dominated in this case by short-distance contributions. For light nuclei, $\not P\not T$ corrections to the wavefunction now are not only due to $\not P\not T$ pion exchange, but also to short-distance nucleon-nucleon interactions. In general, the increased role of short-distance interactions in the case of χISs reduces the predictive power of our analysis, because of the appearance of more LECs.

Finally, pion physics is suppressed also in the case of $\not{P}\not{T}$ from the qEDM. In this case the suppression comes from the need to integrate out the photon to produce purely hadronic operators, which leads to a factor of $\alpha_{\text{em}}/4\pi$. In this case, EDMs of light nuclei are dominated by the nucleon EDM.

Traditionally, when discussing parity and time-reversal violation between nucleons and pions, three non-derivative pion-nucleon interactions are considered on the same footing:

$$\mathcal{L}_{\not{T},\pi N} = -\frac{\bar{g}_0}{F_\pi}\bar{N}\boldsymbol{\tau}\cdot\boldsymbol{\pi}N - \frac{\bar{g}_1}{F_\pi}\pi_3\bar{N}N - \frac{\bar{g}_2}{F_\pi}\pi_3\bar{N}\tau_3 N. \tag{208}$$

When one takes into account the chiral properties of the fundamental sources of $\not{P}\not{T}$, the $\bar{\theta}$ term and the dimension-six sources in Eq. (54), this picture changes. First, these interactions are accompanied by others with further pion fields. Second, and more important, the three interactions are not of the same size. For example, in the case of the chiral-symmetry breaking but isospin-conserving $\bar{\theta}$ term, at LO only $\bar{g}_0$ appears [4]. The couplings $\bar{g}_1$ and $\bar{g}_2$ are respectively suppressed by two and three powers of m_π/M_{QCD} [32], although, once values from the connection to isospin violation in the quark masses [32] are taken into account, $\bar{g}_0$ is numerically smaller than expected [38]. If one wants to study observables sensitive to $\bar{g}_1$ one needs to take into account the full NNLO Lagrangian constructed above, which includes, apart from $\bar{g}_1$, also derivative pion-nucleon and multi-pion-nucleon interactions. An example of this is the $\not{P}\not{T}$ two-nucleon potential which, at the order where $\bar{g}_1$ appears, has a rich and non-trivial momentum dependence in the isoscalar channel [35].

For qCEDM, arguably the most natural case is the one where the isoscalar and isovector components are of similar size, $|\tilde{\delta}_0| \simeq |\tilde{\delta}_3|$. In this scenario the $\bar{g}_0$ and $\bar{g}_1$ interactions appear at the same order. The third pion-nucleon coupling $\bar{g}_2$ comes in two orders down in the chiral expansion. The case of a dominant isoscalar qCEDM, $|\tilde{\delta}_0| \gg |\tilde{\delta}_3|$, generates an identical low-energy $\not{P}\not{T}$ Lagrangian as the $\bar{\theta}$ term, making it impossible to separate these two scenarios from low-energy $\not{P}\not{T}$ observables alone. To this goal, more input from techniques like lattice QCD is needed. The appearance of a dominant isovector qCEDM, $|\tilde{\delta}_3| \gg |\tilde{\delta}_0|$, implies that $\bar{g}_0$ and $\bar{g}_1$ are approximately of the same order, although the former is expected to be somewhat smaller due to the extra factor ε.

The pattern of non-derivative pion-nucleon interactions in the case of the chiral- and isospin-symmetry-breaking four-quark operator FQLR is very similar to that of a dominant isovector qCEDM. The first difference is the appearance of interactions involving multiple pions, which are hard to isolate. A second difference is the appearance of $\not{P}\not{T}$ mesonic operators which, through loops, add to the $\not{P}\not{T}$ PNFF momentum dependence, however at subleading order. The same mesonic operator induces a LO three-nucleon potential, which could be relevant for nuclei with $A \geq 3$. Regardless, separating the FQLR from an isovector qCEDM with hadronic observables requires very precise measurements and is, at this point, not very likely. The search for T violation in β decay could provide additional clues [22].

The χISs —gCEDM and two four-quark interactions— give rise to a similar hierarchy between the non-derivative pion-nucleon couplings as an isovector qCEDM, although in this case it is $\bar{g}_1$ that is expected to be smaller than $\bar{g}_0$ by a factor ε. The LEC $\bar{g}_2$ is suppressed by only one power of m_π/M_{QCD}. Also, at the same order as $\bar{g}_0$ and $\bar{g}_1$, derivative pion-nucleon interactions appear. Within our framework a separation of the different chiral-invariant operators themselves is not possible. For that more advanced techniques than NDA are required to estimate the size of the LECs.

Table 1: The LO scaling of important $\not{P}\not{T}$ LECs for the different $\not{P}\not{T}$ sources. The $\bar{g}_i$ are the non-derivative πN couplings, $\bar{d}_0$ and $\bar{d}_1$ isoscalar and isovector short-range nucleon EDMs, $\bar{c}_0$ and $\bar{c}_1$ isoscalar and isovector magnetic $\gamma\pi N$ interactions, and $\bar{C}_{1,2}$ and $\bar{C}_{3,4}$ isoscalar and isovector NN interactions. (The + in some of the entries should not be taken literally but only as an indication that the LECs get contributions from two sources.)

Source	$\bar{\theta}$	qCEDM	FQLR	qEDM	χISs
$\bar{g}_0$	$\bar{\theta}\frac{m_\pi^2}{M_{\mathrm{QCD}}}$	$(\tilde{\delta}_0+\varepsilon\tilde{\delta}_3)\frac{m_\pi^2 M_{\mathrm{QCD}}}{M_{\not{T}}^2}$	$\varepsilon\xi\frac{M_{\mathrm{QCD}}^3}{M_{\not{T}}^2}$	$(\delta_0+\delta_3)\frac{\alpha_{\mathrm{em}}}{4\pi}\frac{m_\pi^2 M_{\mathrm{QCD}}}{M_{\not{T}}^2}$	$w\frac{m_\pi^2 M_{\mathrm{QCD}}}{M_{\not{T}}^2}$
$\bar{g}_1/\bar{g}_0$	$\varepsilon\frac{m_\pi^2}{M_{\mathrm{QCD}}^2}$	$\frac{\tilde{\delta}_3}{\tilde{\delta}_0+\varepsilon\tilde{\delta}_3}$	$\frac{1}{\varepsilon}$	1	ε
$\bar{g}_2/\bar{g}_0$	$\frac{\alpha_{\mathrm{em}}}{4\pi}$	$\frac{\varepsilon\tilde{\delta}_3}{\tilde{\delta}_0+\varepsilon\tilde{\delta}_3}\frac{m_\pi^2}{M_{\mathrm{QCD}}^2}$	$\varepsilon\frac{m_\pi^2}{M_{\mathrm{QCD}}^2}$	$\frac{\delta_3}{\delta_0+\delta_3}$	$\frac{\alpha_{\mathrm{em}}}{4\pi}\frac{M_{\mathrm{QCD}}^2}{m_\pi^2}$
$\bar{d}_0$	$e\bar{\theta}\frac{m_\pi^2}{M_{\mathrm{QCD}}^3}$	$e(\tilde{\delta}_0+\tilde{\delta}_3)\frac{m_\pi^2}{M_{\not{T}}^2 M_{\mathrm{QCD}}}$	$e\xi\frac{M_{\mathrm{QCD}}}{M_{\not{T}}^2}$	$e\delta_0\frac{m_\pi^2}{M_{\not{T}}^2 M_{\mathrm{QCD}}}$	$ew\frac{M_{\mathrm{QCD}}}{M_{\not{T}}^2}$
$\bar{d}_1/\bar{d}_0$	1	1	1	$\frac{\delta_3}{\delta_0}$	1
$\bar{c}_0/\bar{d}_0$	1	1	1	1	$\frac{m_\pi^2}{M_{\mathrm{QCD}}^2}$
$\bar{c}_1/\bar{d}_0$	1	1	1	$\frac{\delta_3}{\delta_0}$	$\frac{m_\pi^2}{M_{\mathrm{QCD}}^2}$
$\bar{C}_{1,2}$	$\bar{\theta}\frac{m_\pi^2}{F_\pi^2 M_{\mathrm{QCD}}^3}$	$(\tilde{\delta}_0+\varepsilon\tilde{\delta}_3)\frac{m_\pi^2}{F_\pi^2 M_{\not{T}}^2 M_{\mathrm{QCD}}}$	$\xi\frac{M_{\mathrm{QCD}}}{F_\pi^2 M_{\not{T}}^2}$	$(\delta_0+\tilde{\delta}_3)\frac{\alpha_{\mathrm{em}}}{4\pi}\frac{m_\pi^2}{F_\pi^2 M_{\not{T}}^2 M_{\mathrm{QCD}}}$	$w\frac{M_{\mathrm{QCD}}}{F_\pi^2 M_{\not{T}}^2}$
$\bar{C}_{3,4}/\bar{C}_{1,2}$	$\varepsilon\frac{m_\pi^2}{M_{\mathrm{QCD}}^2}$	$\frac{\tilde{\delta}_3}{\tilde{\delta}_0+\varepsilon\tilde{\delta}_3}$	$\frac{1}{\varepsilon}$	1	$\varepsilon\frac{m_\pi^2}{M_{\mathrm{QCD}}^2}$

Only for qEDM are the three non-derivative pion-nucleon couplings expected to be of the same order. But in this case all these couplings are pretty unimportant. The scalings of these couplings for all sources are summarized in the first three rows of Table 1.

The fact that different sources of $\not{P}\not{T}$ are responsible for different hierarchies between the non-derivative $\not{P}\not{T}$ couplings has important implications for the $\not{P}\not{T}$ electromagnetic moments of nuclei, of which the best example is the deuteron EDM. In calculations where the non-derivative pion-nucleon interactions are assumed to be equally sized, it is found that the deuteron EDM is dominated by $\bar{g}_1$ and is larger than the nucleon EDM [62]. The MQM gets contributions of similar size from $\bar{g}_0$ and $\bar{g}_1$. Similarly, it is found that for the helion EDM both $\bar{g}_0$ and $\bar{g}_1$ are important [67]. Taking into account the chiral properties of the fundamental $\not{P}\not{T}$ sources, these conclusions only hold for the isovector qCEDM and FQLR, since for these sources $\bar{g}_0$ and $\bar{g}_1$ interactions appear at leading order. More generally, we see no particular reason to insist on including $\bar{g}_2$ for any source, at least in calculations of light systems.

Instead, what our analysis shows is that one should include other types of $\not{P}\not{T}$ operators, such as nucleon-photon, pion-nucleon-photon, and nucleon-nucleon interactions. In Ref. [36], we argued that the EDMs of light nuclei depend on six different LECs. In addition to the two $\not{P}\not{T}$ pion-nucleon interactions $\bar{g}_0$ and $\bar{g}_1$, they consist of the short-range isoscalar and isovector nucleon EDMs $\bar{d}_0$ and $\bar{d}_1$, and the short-range isoscalar $\not{P}\not{T}$ nucleon-nucleon interactions $\bar{C}_0$ and $\bar{C}_1$. In Ref. [36] the FQLR source was not considered. As we saw above, the surviving three-pion vertex introduces a dependence of observables on $\bar{\Delta}_{\rm LR}^{(-4)}$. However, $\bar{\Delta}_{\rm LR}^{(-4)}$ can be eliminated in favor of $\bar{g}_0$ and CP-even LECs via Eq. (122), so we can still consider LO nuclear EDMs to be expressed in terms of those six LECs. Higher electromagnetic moments, such as the deuteron MQM [43, 37], can depend on additional interactions such as the isoscalar $\not{P}\not{T}$ pion-nucleon-photon vertex $\bar{c}_0$. We summarize the size of these important LECs (extended with some others) in Table 1.

The Lagrangians derived in this work for each of the dimension-six $\not{P}\not{T}$ sources serve as the basis for the calculations of hadronic and nuclear observables. In fact, the $\not{P}\not{T}$ moments of the nucleon [29, 43, 33] and of light nuclei [34, 36, 37, 38] have already been calculated with some of the interactions constructed here and, for the theta term, in Ref. [32]. Other observables such as the EDMs and Schiff moments of heavier nuclei can now be tackled with the same method.

Acknowledgements

We thank D. Boer, J. Bsaisou, C. Hanhart, C.-P. Liu, M. Ramsey-Musolf, I. Stetcu, and A. Wirzba for helpful discussions. J. de Vries and E. Mereghetti acknowledge discussions with W. den Dunnen and W. Dekens, and with H. Murayama, respectively. U. van Kolck acknowledges discussions with W. Hockings at very early stages of this work, and the hospitality of the KVI Groningen on many occasions. This research was supported by the Dutch Stichting FOM under programs 104 and 114 (JdV, RGET) and by the US DOE under grants DE-FG02-06ER41449 (EM) and DE-FG02-04ER41338 (EM, UvK).

References

[1] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).

[2] C. Jarlskog, Phys. Rev. Lett. **55**, 1039 (1985).

[3] G. 't Hooft, Phys. Rev. Lett. **37**, 8 (1976); Phys. Rev. D **14**, 3432 (1976); **18**, 2199(E) (1978).

[4] R. J. Crewther, P. Di Vecchia, G. Veneziano, and E. Witten, Phys. Lett. B **88**, 123 (1979); **91**, 487(E) (1980).

[5] A. Riotto and M. Trodden, Ann. Rev. Nucl. Part. Sci. **49**, 35 (1999).

[6] I. B. Khriplovich and S. K. Lamoreaux, *CP Violation Without Strangeness: Electric Dipole Moments of Particles, Atoms, and Molecules* (Springer Verlag, Berlin, 1997).

[7] M. Pospelov and A. Ritz, Ann. Phys. **318**, 119 (2005).

[8] B. H. J. McKellar, S. R. Choudhury, X. G. He, and S. Pakvasa, Phys. Lett. B **197**, 556 (1987).

[9] T. M. Ito, J. Phys. Conf. Ser. **69**, 012037 (2007), nucl-ex/0702024; K. Bodek *et al.*, arXiv:0806.4837 [nucl-ex].

[10] C. A. Baker *et al.*, Phys. Rev. Lett. **97**, 131801 (2006).

[11] W. C. Griffiths *et al.*, Phys. Rev. Lett. **102**, 101601 (2009).

[12] F. J. M. Farley *et al.*, Phys. Rev. Lett. **93**, 052001 (2004); Y. F. Orlov, W. M. Morse, and Y. K. Semertzidis, Phys. Rev. Lett. **96**, 214802 (2006); C. J. G. Onderwater, J. Phys. Conf. Ser. **295**, 012008 (2011).

[13] S. Weinberg, Phys. Rev. Lett. **43**, 1566 (1979).

[14] W. Buchmüller and D. Wyler, Nucl. Phys. B **268**, 621 (1986).

[15] A. De Rújula, M. B. Gavela, O. Pène, and F. J. Vegas, Nucl. Phys. B **357**, 311 (1991).

[16] M. J. Ramsey-Musolf and S. Su, Phys. Rept. **456**, 1 (2008).

[17] B. Grzadkowski, M. Iskrzynski, M. Misiak, and J. Rosiek, JHEP **1010**, 085 (2010).

[18] J. Ellis, S. Ferrara, and D. V. Nanopoulos, Phys. Lett. B **114**, 231 (1982); J. Polchinski and M. B. Wise, Phys. Lett. B **125**, 393 (1983); M. Dugan, B. Grinstein, and L. Hall, Nucl. Phys. B **255**, 413 (1985).

[19] R. Arnowitt, J. Lopez, and D. Nanopoulos, Phys. Rev. D **42**, 2423 (1990); R. Arnowitt, M. Duff, and K. Stelle, Phys. Rev. D **43**, 3085 (1991); Y. Kizuri and N. Oshimo, Phys. Rev. D **45**, 1806 (1992).

[20] S. Weinberg, Phys. Rev. Lett. **63**, 2333 (1989).

[21] E. Braaten, C. S. Li, and T. C. Yuan, Phys. Rev. Lett. **64**, 1709 (1990).

[22] J. Ng and S. Tulin, Phys. Rev. D **85**, 033001 (2012).

[23] S. Weinberg, Phys. Lett. B **251**, 288 (1990); Nucl. Phys. B **363**, 3 (1991).

[24] S. Weinberg, Physica **96A**, 327 (1979); J. Gasser and H. Leutwyler, Ann. Phys. **158**, 142 (1984); Nucl. Phys. B **250**, 465 (1985).

[25] S. Weinberg, *The Quantum Theory of Fields*, Vol. 2 (Cambridge University Press, Cambridge, 1996).

[26] U.-G. Meißner, arXiv:hep-ph/9711365.

[27] P. F. Bedaque and U. van Kolck, Ann. Rev. Nucl. Part. Sci. **52**, 339 (2002).

[28] S. D. Thomas, Phys. Rev. D **51**, 3955 (1995).

[29] W. H. Hockings and U. van Kolck, Phys. Lett. B **605**, 273 (2005).

[30] W. H. Hockings, Ph.D. dissertation, University of Arizona (2006); E. Mereghetti, Ph.D. dissertation, University of Arizona (2011); J. de Vries, Ph.D. dissertation, University of Groningen (2012).

[31] S. Narison, Phys. Lett. B **666**, 455 (2008).

[32] E. Mereghetti, W. H. Hockings, and U. van Kolck, Ann. Phys. **325**, 2363 (2010).

[33] E. Mereghetti, J. de Vries, W.H. Hockings, C.M. Maekawa, and U. van Kolck, Phys. Lett. B **696**, 97 (2011).

[34] J. de Vries, E. Mereghetti, R. G. E. Timmermans, and U. van Kolck, Phys. Rev. Lett. **107**, 091804 (2011).

[35] C. M. Maekawa, E. Mereghetti, J. de Vries, and U. van Kolck, Nucl. Phys. A **872**, 117 (2011).

[36] J. de Vries, R. Higa, C.-P. Liu, E. Mereghetti, I. Stetcu, R. G. E. Timmermans, and U. van Kolck, Phys. Rev. C **84**, 065501 (2011).

[37] C.-P. Liu, J. de Vries, E. Mereghetti, R. G. E. Timmermans, and U. van Kolck, Phys. Lett. B **713** (2012) 447.

[38] J. Bsaisou, C. Hanhart, S. Liebig, U.-G. Meißner, A. Nogga, and A. Wirzba, arXiv:1209.6306 [hep-ph].

[39] H.-Y. Cheng, Phys. Rev. D **44**, 166 (1991); A. Pich and E. de Rafael, Nucl. Phys. B **367**, 313 (1991); P. Cho, Phys. Rev. D **48**, 3304 (1993); B. Borasoy, Phys. Rev. D **61**, 114017 (2000); F.-K. Guo and U.-G. Meißner, arXiv:1210.5887 [hep-ph].

[40] K. Ottnad, B. Kubis, U.-G. Meißner, and F.-K. Guo, Phys. Lett. B **687**, 42 (2010).

[41] E. E. Jenkins and A. V. Manohar, Phys. Lett. B **255**, 558 (1991).

[42] M. E. Luke and A. V. Manohar, Phys. Lett. B **286**, 348 (1992); W. Kilian and T. Ohl, Phys. Rev. D **50**, 4649 (1994); R. Sundrum, Phys. Rev. D **59**, 085009 (1999); J. de Vries, E. Mereghetti, R.G.E. Timmermans, and U. van Kolck, in preparation.

[43] J. de Vries, E. Mereghetti, R.G.E. Timmermans, and U. van Kolck, Phys. Lett. B **695**, 268 (2011).

[44] V. Baluni, Phys. Rev. D **19**, 2227 (1979).

[45] K. Fujikawa, Phys. Rev. Lett. **42**, 1195 (1979).

[46] G. Isidori, J. F. Kamenik, Z. Ligeti, and G. Perez, Phys. Lett. B **711**, 46 (2012).

[47] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **108**, 111602 (2012).

[48] H. An, X. Ji, and F. Xu, JHEP **1002**, 043 (2010).

[49] A. Cordero-Cid, J. M. Hernandez, G. Tavares-Velasco, and J. J. Toscano, J. Phys. G **35**, 025004 (2008); J. F. Kamenik, M. Papucci, and A. Weiler, Phys. Rev. D **85**, 071501 (2012).

[50] A. De Rújula, M. B. Gavela, O. Pène, and F. J. Vegas, Phys. Lett. B **245**, 640 (1990); G. Degrassi, E. Franco, S. Marchetti, and L. Silvestrini, JHEP **0511**, 044 (2005); J. Hisano, K. Tsumura, and M. J. S. Yang, Phys. Lett. B **713**, 473 (2012).

[51] F. Xu, H. An, and X. Ji, JHEP **1003**, 088 (2010).

[52] M. E. Pospelov, Phys. Lett. B **328**, 441 (1994); A. Czarnecki and B. Krause, Phys. Rev. Lett. **78**, 4339 (1997).

[53] T. Ibrahim and P. Nath, Rev. Mod. Phys. **80**, 577 (2008).

[54] U. van Kolck, Ph.D. dissertation, University of Texas (1993); Few Body Syst. Suppl. **9**, 444 (1995).

[55] A. V. Manohar and H. Georgi, Nucl. Phys. B **234**, 189 (1984).

[56] J. L. Friar, U. van Kolck, M. C. M. Rentmeester, and R. G. E. Timmermans, Phys. Rev. C **70**, 044001 (2004).

[57] S. Weinberg, Trans. N. Y. Acad. Sci. **38**, 185 (1977).

[58] W. Fischler, S. Paban, and S. Thomas, Phys. Lett. B **289**, 373 (1992).

[59] O. Lebedev, K. A. Olive, M. Pospelov, and A. Ritz, Phys. Rev. D **70**, 016003 (2004).

[60] S. R. Beane, K. Orginos, and M. J. Savage, Nucl. Phys. B **768**, 38 (2007).

[61] J. Gasser, H. Leutwyler, and M. E. Sainio, Phys. Lett. B **253**, 252 (1991); J. M. Alarcon, J. Martin Camalich, and J. A. Oller, Phys. Rev. D **85**, 051503 (2012); G. S. Bali *et al.* Nucl. Phys. B **866**, 1 (2013).

[62] I. B. Khriplovich and R. A. Korkin, Nucl. Phys. A **665**, 365 (2000); C.-P. Liu and R. G. E. Timmermans, Phys. Rev. C **70**, 055501 (2004); Phys. Lett. B **634**, 488 (2006).

[63] C.-H. Seng, H. Patel, J. de Vries, E. Mereghetti *et al.*, in progress.

[64] S. R. Beane, P. F. Bedaque, M. J. Savage, and U. van Kolck, Nucl. Phys. A **700**, 377 (2002); A. Nogga, R. G. E. Timmermans, and U. van Kolck, Phys. Rev. C **72**, 054006 (2005); M. Birse, Phys. Rev. C **74**, 014003 (2006); M. Pavón Valderrama, Phys. Rev. C **83**, 024003 (2011); **84**, 064002 (2011); B. Long and C.J. Yang, Phys. Rev. C **84**, 057001 (2011); **85**, 034002 (2012); **86**, 024001 (2012).

[65] J. L. Friar, Few Body Syst. **22**, 161 (1997).

[66] U. van Kolck, Phys. Rev. C **49**, 2932 (1994).

[67] I. Stetcu, C.-P. Liu, J. L. Friar, A. C. Hayes, and P. Navrátil, Phys. Lett. B **665**, 168 (2008); Y.-H. Song, R. Lazauskas, and V. Gudkov, arXiv:1211.3762 [nucl-th].